



Canadian Council of Ministers
of the Environment Le Conseil canadien
des ministres
de l'environnement

**Scientific Criteria Document
for the Development of the
Canadian Water Quality Guidelines for**

BORON

PN 1437

ISBN 978-1-896997-89-6 PDF

© Canadian Council of Ministers of the Environment, 2009

PREFACE

CCME has recently updated the 1991 protocol for the derivation of Water Quality Guidelines for the Protection of Aquatic Life (WQGPAqL) and developed a tiered approach to guideline derivation, which now includes the use of a statistical extrapolation method in addition to the original assessment factor approach. This updated protocol has been peer reviewed and adopted as the new tiered approach. The first tier of the CCME (2007) national approach is to make use of species sensitivity distributions (SSDs). This approach uses all available toxicity data for species to derive the final guideline as the data is fitted to a specified model or distribution, from which is calculated an exposure concentration that is to be protective of a specified percentage of species (e.g. 95%). If the data is insufficient to model an SSD curve, then a second or third tier guideline would be developed using an assessment factor method approach (dividing the lowest toxicity value from a high quality study by an uncertainty or safety factor), which is similar to the 1991 CCME guideline development protocol.

The 2007 CCME protocol for the derivation of WQGPAqL now provides guidance on setting both a short-term benchmark concentration and a long-term guideline. Short-term exposure benchmark concentrations are meant to protect only a specified fraction of individuals for a defined short-term exposure period from severe effects such as lethality or immobilization, as a result of spill events to aquatic-receiving environments or from infrequent releases of short-lived/non-persistent substances (CCME, 2007). In contrast, long-term exposure guidelines adhere to the CCME guiding principle and are meant to protect all forms of aquatic life (all species, all life stages) for indefinite exposure periods (CCME, 2007). There are three tiers in place for the setting of CCME (2007) WQGPAqL. **Type A** guidelines are derived using a species sensitivity distribution (SSD) approach when there are adequate primary and secondary toxicity data to satisfactorily fit a SSD curve. **Type B** guidelines are derived for substances that either have inadequate or insufficient toxicity data for the SSD approach (i.e., Type A guideline), but for which enough toxicity data from a minimum number of primary and/or secondary studies are available. The Type B guideline approach is further divided into the **Type B1** and **Type B2** guidelines, based upon the quantity and quality of available toxicity data.

In this report, a Type A short-term boron benchmark concentration and a long-term boron water quality guideline for the protection of aquatic life have been developed using the national species sensitivity distribution method (CCME, 2007).

ACKNOWLEDGEMENTS

This Scientific Criteria Document has been contributed to CCME by the Standards Development Branch, Ontario Ministry of Environment.

An original draft of this document was prepared by Nicole Knezevich and Ruth Hull (Intrinsic Environmental Sciences Inc., Mississauga, ON) as well as Dwayne Moore and Scott Teed (Intrinsic Environmental Sciences Inc, Ottawa, ON). Additional work and completion of the document was done by Monica Nowierski and Tim Fletcher (Ontario Ministry of the Environment, Standards Development Branch). Thanks are extended to the following individuals for their help in the preparation of the boron Water Quality Guideline Supporting Document:

- David Poirier and Jordana VanGeest, from the Laboratory Services Branch, Ontario Ministry of the Environment, for conducting boron toxicity test
- Peer reviewers from the CCME (Canadian Council of Ministers of the Environment) Water Quality Task Group for their comments: Marc Demers, Sylvie Cloutier, Isabelle Guay, and Kelly Potter.
- Comments were also received from Cara Clairman (Ontario Power Generation) and Dr. Michael C Harrass (Rio Tinto Minerals).
- Marc Demers (National Guidelines Standards Office) for the addition of data for the generation of the boron species sensitivity distributions.

Table of Contents

ACKNOWLEDGEMENTS	iii
SUMMARY	2
RÉSUMÉ	3
1.0 INTRODUCTION	4
1.1 <i>Production and Uses</i>	4
1.2 <i>Aquatic Sources, Properties and Fate</i>	4
1.3 <i>Ambient Concentrations in Canadian Waters</i>	8
1.3 <i>Ambient Concentrations in Canadian Waters</i>	9
2.0 TOXICITY TO AQUATIC ORGANISMS	11
2.1 <i>Acute Toxicity</i>	12
2.1.1 <i>Vertebrates</i>	12
2.1.2 <i>Invertebrates</i>	12
2.2 <i>Chronic Toxicity</i>	13
2.2.1 <i>Vertebrates</i>	13
2.2.2 <i>Invertebrates</i>	15
2.2.3 <i>Other Organisms (Algae, Macrophytes, etc.)</i>	16
2.3 <i>Summary of Toxicity Data</i>	16
2.4 <i>Effects of Water Quality Parameters on Toxicity</i>	18
2.5 <i>Essentiality and Deficiency</i>	<i>Error! Bookmark not defined.</i>
3.0 BIOACCUMULATION	20
4.0 IMPACT ON TASTE AND ODOUR OF WATER AND FISH TAINING	21
5.0 MUTAGENICITY	21
6.0 DERMAL AND OTHER EFFECTS	22
7.0 METHOD DETECTION LIMITS	22
8.0 DERIVATION OF SPECIES SENSITIVITY DISTRIBUTIONS	22
8.1 <i>Short-term SSD Derivation</i>	23
8.2 <i>Long-term SSD Derivation</i>	27
9.0 RESEARCH NEEDS	34
10.0 OBJECTIVES OF OTHER AGENCIES	34
11.0 REFERENCES	36

List of Tables

Table 1- 1 Physical - Chemical Properties Boron	6
Table 1- 2 Physical - Chemical Properties Boron	7
Table 1- 3 Physical - Chemical Properties	8
Table 1- 4 Boron Ontario Surface Water Concentrations (mg/L) ^a	9
Table 1- 5 Historical (1983 to 1985) Boron Surface Water Concentrations (mg/L) in Lake Superior and Lake Ontario	10
Table 1- 6 Boron Surface Water Concentrations in Canada (mg/L)	11
Table 8- 1 Minimum Data Set Requirements for the Gereneration of a Short-Term Freshwater Benchmark Concentration	24
Table 8- 2 Short-term LC ₅₀ s for species exposed to boron in freshwater	24
Table 8- 3 Studies Used to Derive Geometric Means for Short-Term Data	25
Table 8- 4 Short-term Concentration for Boron Resulting from the SSD Method	26
Table 8- 5 Minimum Data Set Requirements for the Generation of a long-term freshwater CWQG	27
Table 8- 6 Long-term low effects and no effects concentrations for species exposed to boron in freshwater	29
Table 8- 7 Studies Used to Derive Geometric Means for Long-Term Data	30
Table 8- 8 Long-term CWQG for Boron Resulting from the SSD Method	32

List of Figures

Figure 1 The Effects of Boric Acid Administered to Embryo-Larval Stages of Trout in Soft Water (Birge and Black, 1977)	15
Figure 2 Toxicity ranges exhibited by aquatic organisms exposed to boron.	15
Figure 3 SSD for boron in freshwater derived by fitting the log-Gompertz model to the logarithm of acceptable short-term LC ₅₀ s of thirteen aquatic species versus Hazen plotting position (proportion of species affected).	27
Figure 4 SSD for boron in freshwater derived by fitting the log-Normal model to the logarithm of acceptable long-term no- and low-effect endpoints for twenty-eight aquatic species versus Hazen plotting position (proportion of species affected)	33

SUMMARY

CCME Water Quality Guidelines for the Protection of Aquatic Life (CWQGs PAL) abide by the following general guiding principle: to protect all forms of aquatic life and all aspects of aquatic life cycles. The clear intention is to protect all species during all life stages during indefinite exposure to the water. CCME has recently updated the 1991 protocol for the derivation of CWQGs PAL and developed a tiered approach to guideline derivation, which now includes the use of species sensitivity distributions in addition to the original assessment factor approach. In this report, boron water quality guidelines for the protection of aquatic life have been developed using the national species sensitivity distribution method (or SSD Type A guideline derivation method). Both a short-term exposure benchmark concentrations (29 mg/L) and a long-term (1.5 mg/L) exposure guideline have been developed. Short-term exposure benchmark concentrations are derived using severe effects data (such as lethality) of defined short-term exposure periods (24-96 h). These benchmark concentrations identify estimators of severe effects to the aquatic ecosystem and are intended to give guidance on the impacts of severe, but transient, situations (e.g., spill events to aquatic receiving environments and infrequent releases of short-lived/nonpersistent substances). Short-term benchmark concentrations *do not* provide guidance on protective levels of a substance in the aquatic environment, as short-term benchmark concentrations are levels which *do not* protect against adverse effects. Long-term exposure guidelines are intended to protect all forms of aquatic life for indefinite exposure periods (≥ 7 d exposures for fish and invertebrates, ≥ 24 h exposures for aquatic plants and algae).

Canadian Water Quality Guideline (CWQG) for Boron (mg L⁻¹) for the Protection of Aquatic Life		
	Long-Term Exposure Guideline	Short-Term Exposure Benchmark Concentration
Freshwater	1.5	29
Marine	NRG	NRG

NRG = No recommended guideline

RÉSUMÉ

Les Recommandations canadiennes pour la qualité des eaux en vue de protéger la vie aquatique (RCQE-PVA) reposent sur le même principe directeur général, soit protéger toutes les formes de vie aquatique et tous les aspects des cycles biologiques en milieu aquatique. Elles visent à protéger toutes les espèces à tous les stades de vie pendant des périodes d'exposition indéterminées dans le milieu aquatique. Le CCME a récemment mis à jour le protocole d'élaboration des RCQE-PVA de 1991 et a développé une démarche en plusieurs étapes qui tient compte de la distribution de la sensibilité des espèces (DSE) en plus des facteurs d'évaluation déjà utilisés. Dans le présent rapport, les recommandations relatives au bore ont été élaborées à l'aide de la méthode nationale de la DSE (méthode d'élaboration de recommandations de type A). Une recommandation pour une exposition de courte durée (29 mg/L) et une recommandation pour une exposition de longue durée (1,5 mg/L) ont été élaborées. Les recommandations pour une exposition de courte durée sont élaborées au moyen de données sur les effets graves (comme la létalité) associés à des périodes d'exposition de courte durée bien définies (de 24 à 96 h). Elles donnent une indication des concentrations pouvant entraîner des effets graves pour les écosystèmes aquatiques et renseignent sur les impacts d'événements graves, mais transitoires (p. ex. déversements dans le milieu aquatique et rejets peu fréquents de substances non persistantes ou à courte durée de vie). Elles *ne donnent pas* d'indication sur les concentrations qui assurent la protection des organismes aquatiques et *ne protègent pas* contre les effets nocifs des substances. Les recommandations pour une exposition de longue durée visent à protéger toutes les formes de vie aquatique pendant des périodes d'exposition indéterminées (≥ 7 jours pour les poissons et les invertébrés et ≥ 24 h pour les végétaux aquatiques et les algues).

Recommandations canadiennes pour la qualité des eaux (RCQE) en vue de protéger la vie aquatique – bore (mg L⁻¹)		
	Exposition de longue durée	Exposition de courte durée
Eaux douces	1.5	29
Eaux marines	AR	AR

AR = aucune recommandation

1.0 INTRODUCTION

1.1 Production and Uses

Boron is ubiquitous in the environment, occurring naturally in over 80 minerals and constituting 0.001% of the Earth's crust (U.S. EPA, 1987). The highest concentrations of boron are found in sediments and sedimentary rock, particularly in clay rich marine sediments. The high boron concentration in seawater (4.5 mg B/L), ensures that marine clays are rich in boron relative to other rock types (Butterwick *et al.*, 1989). The most significant source of boron is seasalt aerosols, where annual input of boron to the atmosphere is estimated to be 1.44 Tg B/year, where Tg = 10^{12} g (Park and Schlesinger, 2002). In addition to atmospheric deposition, boron is also released into the environment very slowly at low concentrations by natural weathering processes. Due to the extensive occurrence of clay-rich sedimentary rocks on the Earth's land surfaces, the majority of boron mobilized into soils and the aquatic environment by weathering probably stems from this source. Natural weathering is estimated to release more boron into the environment worldwide than industrial sources (Butterwick *et al.*, 1989) as boron is tied up in many industrial products such as glass. Natural weathering (chemical and mechanical) of minerals from carbonate rocks has been estimated to mobilize 0.193 Tg B/year (Park and Schlesinger, 2002). Other natural sources include volcanic emissions (0.017-0.022 Tg B/yr), soil dust (0.017-0.033 Tg B/yr), and plant aerosols (0.004 Tg B/yr). Volcanic emissions release boric acid and boron trifluoride; therefore, the concentrations of boron in water in volcanic regions are high (Health Canada, 1990). The evaporation of sea water from closed basins is a commercial source of boron (Durocher, 1969). The estimated amounts of boron introduced into the atmosphere as a result of fossil fuel combustion, biomass burning and other human activities (e.g. manufacturing and incineration) are 0.20, 0.26-0.43, and 0.07 Tg B/yr, respectively (Park and Schlesinger, 2002).

Boron is produced anthropogenically by the chemical reduction of boron compounds with reactive metals, either by non-aqueous electrolytic reduction or thermal decomposition. Highly purified boron is produced by zone-refining or other thermal techniques (Stokinger, 1981; U.S. Bureau of Mines, 1989). Borax, found in playa (intermittent) lakes and other evaporite deposits, is used to produce refined sodium borate compounds and boric acid (ATSDR, 1992).

Borates and boric acids are used in glass manufacturing (fiberglass, borosilicate glass, enamel, frit, and glaze), soaps and detergents, flame retardants, and neutron absorbers for nuclear installations (WHO, 2003). Boric acid, borates and perborates have been used in mild antiseptics, cosmetics, pharmaceuticals, as antioxidants for soldering, cleaning products/detergents, boron neutron capture therapy and agricultural fertilizers (WHO, 2003). Boron compounds are also used in the leather, textile, paint and wood-processing industries (Health Canada, 1990). Borax (or sodium tetraborate) and boric acid are used as insecticides in Canada (Health Canada, 1990). Borax is also used extensively as a cleaning agent and an antimicrobial agent (Health Canada, 1990).

1.2 Aquatic Sources, Properties and Fate

The wet deposition of boron over the continents is estimated to be approximately 0.50 Tg B/yr, where rainwater from continental sites contains less boron when compared to boron from coastal and marine sites (Park and Schlesinger, 2002). Natural weathering (chemical and mechanical) of boron-containing rocks is a major source of boron compounds in water (Butterwick *et al.*, 1989)

and on land (Park and Schlesinger, 2002). The amount of boron released into the aquatic environment varies greatly depending on the surrounding geology. Boron compounds also are released to water in municipal sewage and in waste waters from coal-burning power plants, irrigation, copper smelters and industries using boron (ATSDR, 1992; Howe, 1998). The boron content of wastewater discharges can be significant as borate compounds are present in domestic washing agents (WHO, 2003). With respect to Canadian wastewater discharges, a literature review was conducted to characterize the state of knowledge of municipal effluent (Hydromantis Inc., 2005). The study reported that boron releases from two Western Canadian wastewater treatment plants, releasing either raw or primary treated effluent, ranged from 110 to 180 $\mu\text{g/L}$ (or 0.11 to 0.18 mg/L). Therefore, the study indicates that municipal wastewater effluent in Canada is less likely to be a source, and boron concentrations in water are more likely dependant on the leaching of boron from the surrounding geology (WHO, 2003).

The physical and chemical properties of boron and its aqueous forms are shown in Tables 1-1, 1-2, and 1-3. Elemental boron is insoluble and inert in aqueous solutions. Boron compounds rapidly transform to borates, the naturally occurring forms of boron, when exposed to water. However, no further degradation is possible. The only significant mechanism expected to influence the fate of boron in water is adsorption-desorption reactions with soil and sediment (Rai *et al.*, 1986). The extent of boron adsorption depends on the pH of the water and concentration of boron in solution. The greatest adsorption is observed between a pH of 7.5 and 9.0 (WHO, 2003).

In natural waters, boron forms stable species and exists primarily as undissociated boric acid [$\text{B}(\text{OH})_3$] and complex polyanions (*e.g.*, $\text{B}(\text{OH})_4^-$) (Health Canada, 1990; Howe, 1998; WHO, 2003). These forms of boron are highly soluble and not easily removed from solution by natural mechanisms. Borate and boric acid are in equilibrium depending on the pH of the water. At an acidic pH, boron exists in solution mainly as undissociated boric acid, whereas at alkaline pH it is present as borate ions (Howe, 1998).

Table 1- 1 Physical - Chemical Properties Boron

Compound: Boron		Chemical Formula: B	CAS No: 7440-42-8
Properties			
Molecular Weight (MW):	10.81 g/mol	Budavari <i>et al.</i> , 1989; Weast, 1985; Lide, 2000; Clayton and Clayton, 1982; Sax, 1984; Windholz, 1983; Moss and Nagpal, 2003	
Melting Point:	2,075 °C (Lide, 2000) - 2,300 °C (Weast, 1985; U.S. EPA, 1975)		
Boiling Point:	2,550 (Weast, 1985) - 4,000 (Lide, 2000)		
Physical State at Standard Temperature and Pressure:	Solid	Weast, 1985; WHO, 2003; Moss and Nagpal, 2003	
Dissociation Constant:	no data		
Liquid Density (D):	2.34 g/cm ³ at 20 °C	Weast, 1985; U.S. EPA, 1975	
Molar Volume (MW/D):	4.62 cm ³ /mol (calc.)		
Vapour Pressure (Pa):	1.56x10 ⁻⁵ atm at 2,140 °C	Windholz, 1983; O'Neil, 2001; Budavari <i>et al.</i> , 1989	
Water Solubility:	Insoluble	Weast, 1985; Windholz, 1983; Clayton and Clayton, 1982; Hawley, 1981; Budavari <i>et al.</i> , 1989; U.S. EPA, 1975	
Henry's Law Constant (Ps/Cs):	no data		
Persistence			
B.O.D. (mg/L)	no data		
Breakdown Products:	Boron does not degrade in the environment (ATSDR, 1992)		
Half Life (Days):	no data		
Octanol-Water Partition Co-efficient (Kow)			
Range of Available Kow Values:	Adsorption constants for inorganic constituents cannot be predicted <i>a priori</i> (ATSDR, 1992)		
Final Chosen Log Kow Value:	no data		

Table 1- 2 Physical - Chemical Properties Boron

Compound: Boric acid	Chemical Formula: H_3O_3	CAS No:010043-35-3
Properties		
Molecular Weight (MW):	61.833 (Lide, 2000) - 69.92 g/mol (Weast, 1985)	
Melting Point:	169°C (Weast, 1985; U.S. EPA, 1975) - 17°C (Budavari <i>et al.</i> , 1989; Lide, 2000)	
Boiling Point:	300°C	Weast, 1985; U.S. EPA, 1975
Physical State at Standard Temperature and Pressure:	Solid	Weast, 1985; CHEMINFO, 1996
Dissociation Constant:	5.8×10^{-10} at 25°C (pKa=9.24)	Kirk-Othmer, 1992
Liquid Density (D):	1.435 g/cm ³ at 15°C	Lewis, 1999
	2.46 g/cm ³ at 20°C	Weast, 1985
Molar Volume (MW/D):	no data	
Vapour Pressure (Pa):	negligible at 20°C	Bingham <i>et al.</i> , 2001
Water Solubility:	47.2 g/L at 25°C	Kirk-Othmer, 1992
	50 g/L at 25°C	Shiu <i>et al.</i> , 1990
	63.5 g/L at 30°C	Weast, 1985; U.S. EPA, 1975
	276 g/L at 100°C	U.S. EPA, 1975
Henry's Law Constant (Ps/Cs):	no data	
Persistence		
B.O.D. (mg/L)	no data	
Breakdown Products:	No biotransformation processes have been reported for boron compounds (O'Neil, 2001)	
Half Life (Days):	no data	
Octanol-Water Partition Co-efficient (Kow)		
Range of Available Kow Values:	no data	
Final Chosen Log Kow Value:	no data	

Table 1- 3 Physical - Chemical Properties

Compound: Borax	Chemical Formula: Na ₂ B ₄ O ₇	CAS No: 1303-96-4
-----------------	---	-------------------

Properties		
Molecular Weight (MW):	381.37 g/mol	Weast, 1985; NIOSH, 2003; Lide, 2000
Melting Point:	60-75°C	Weast, 1985; U.S. EPA, 1975 (75°C); Lide, 2000 (75°C)
Boiling Point:	320°C	Weast, 1985; U.S. EPA, 1975; NIOSH, 2003
Physical State at Standard Temperature and Pressure:	Solid	NIOSH, 2003
Dissociation Constant:	no data	
Liquid Density (D):	1.73 g/cm ³ at 20°C	Weast, 1985; NIOSH, 2003; Lide, 2000
Molar Volume (MW/D):	no data	
Vapour Pressure (Pa):	0 mmHg (approx)	NIOSH, 2003; Lide, 2000
Water Solubility:	20 g/L at 0°C	Weast, 1985; U.S. EPA, 1975
	59.3 g/L at 25°C	Shiu <i>et al.</i> , 1990
	1,700 g/L at 100°C	U.S. EPA, 1975
Henry's Law Constant (Ps/Cs):	no data	

Persistence		
B.O.D. (mg/L)	no data	
Breakdown Products:	No biotransformation processes have been reported for boron compounds (O'Neil, 2001)	
Half Life (Days):	no data	

1.3 Ambient Concentrations in Canadian Waters

The majority of the Earth's boron is found in the oceans, with an average concentration of 4.5 mg/L (Weast, 1985). The amount of boron in fresh water depends on factors such as the proximity to marine coastal regions, inputs from industrial and municipal effluents and the geochemical nature of the drainage area (Butterwick *et al.*, 1989). Naturally-occurring boron is present in groundwater, primarily as a result of leaching from rocks and soils containing borates and borosilicates (WHO, 2003).

Concentrations of boron in the surface water of Canada and the USA ranged from 0.02 to 360 mg/L (WHO, 1998). High boron concentrations are indicative of boron-rich deposits. Typical boron concentrations were less than 0.1 mg/L, with a 90th percentile concentration of 0.4 mg/L (WHO, 1998). Sekerka and Lechner (1990) measured boron concentrations in surface water samples obtained from various locations in Ontario. Boron concentrations ranged from 0.025 to 0.072 mg/L (Table 1-4). In addition, historical dissolved, particulate and total boron concentrations in Lake Superior (1983) and Lake Ontario (1985) were characterized by Rossmann and Barres (1988) (Table 1-5).

Table 1- 4 Boron Ontario Surface Water Concentrations (mg/L)^a

Sample Location	Mean Boron Concentration (SD) (mg/L)
Burlington Bay	0.0654
Hidden Valley Creek	0.0524
Roseland Creek	0.0303
Shore Acres Creek	0.0379
Appleby Creek	0.0342
Lake Ontario	0.0323 (1.17x10 ⁻³)
Lake Ontario (Bay I)	0.0719 (3.5x10 ⁻⁴)
Lake Ontario (Bay II)	0.0630 (1.92x10 ⁻³)
Lake Ontario	0.0251
Hamilton Bay	0.0593
Northern Ontario bog water	0.0533

^a Obtained from Sekerka and Lechner, 1990

Table 1- 5 Historical (1983 to 1985) Boron Surface Water Concentrations (mg/L) in Lake Superior and Lake Ontario

Boron Form	Boron Concentration (mg/L)	
	Lake Superior	Lake Ontario
Dissolved	0.027	0.047
Particulate	0.0009	0.0055
Total	NA	0.058

NA Not available

The National Water Quality Monitoring Office of Environment Canada provided boron concentration data measured in surface waters from various locations across Canada (C. Lochner 2008, pers. com.). This data is presented in Table 1-6. Boron concentrations ranged from 0.0001-0.951 mg L⁻¹ (extractable) in Nova Scotia, 0.008-0.13 mg L⁻¹ (extractable) and 0-0.607 mg L⁻¹ (total) in Newfoundland, and 0.0001-0.402 mg L⁻¹ (extractable) in New Brunswick. For Ontario, data was provided for the Great Lakes and Great Lake connecting channels. Total boron concentrations ranged from 0.006-0.011 mg L⁻¹ in Lake Superior, 0.004-0.018 mg L⁻¹ in Lake Huron, 0.007-0.011 mg L⁻¹ in Georgian Bay, 0.015-0.031 mg L⁻¹ in Lake Erie and 0.018-0.077 mg L⁻¹ in Lake Ontario. Total boron concentrations measured in the St. Clair River, Niagara River and St. Lawrence River were 0.009-0.021 mg L⁻¹, 0.018-0.032 mg L⁻¹, and 0.02-0.032 mg L⁻¹, respectively. Surface water total boron concentrations ranged from 0.0052-0.271 mg L⁻¹ in Manitoba, 0.0001-2.58 mg L⁻¹ in Saskatchewan, 0.0001-0.082 mg L⁻¹ in Alberta, 0.0001-2.3 mg L⁻¹ in the Northwest Territories, and 0.0001-0.006 mg L⁻¹ in the Yukon.

Table 1- 6 Boron Surface Water Concentrations in Canada (mg/L)

Sample Location	Total Boron Concentration Range	Extractable Boron Concentration Range	Dissolved Boron Concentration Range
Maritimes			
New Brunswick	NA	0.0001 - 0.402	NA
Newfoundland	0 - 0.607	0.008 - 0.13	NA
Nova Scotia	NA	0.0001 - 0.951	NA
Ontario - Great Lakes			
Lake Superior	0.006 - 0.011	NA	NA
Lake Huron	0.004 - 0.018	NA	NA
Georgian Bay	0.007 - 0.011	NA	NA
Lake Erie	0.015 - 0.031	NA	NA
Lake Ontario	0.018 - 0.077	NA	NA
Ontario - Great Lakes Connecting Channels			
St. Clair River	0.009 - 0.021	NA	NA
Niagara River	0.018 - 0.032	0.016 - 0.0245	NA
St. Lawrence River	0.02 - 0.032	0.0229 - 0.0252	NA
Praries			
Manitoba	0.0052 - 0.271	0.01 - 0.11	0.002 - 0.219
Saskatchewan	0.0001 - 2.58	NA	0.0001 - 2.71
Alberta	0.0001 - 0.082	0.01 - 0.13	0.0001 - 0.242
Territories			
Northwest Territories	0.0001 - 2.3	0.01 - 0.16	0.0001 - 0.0568
Yukon Territory	0.0001 - 0.006	0.01	0.0016 - 0.0059

In Quebec, recent data (2004-2006) were obtained in clean conditions, on the acid soluble fraction. Of the 23 rivers sampled, the median boron concentrations ranged from 0.0021 to 0.058 mg/L with a median concentration for all the rivers of 0.0063 mg/L (MDDEP, 2007 unpublished data).

2.0 TOXICITY TO AQUATIC ORGANISMS

Criteria used for classifying available toxicity data as either primary or secondary information are described in “*A Protocol for the Derivation of Water Quality Guidelines for the Protection of Aquatic Life*” (CCME, 2007). The studies from which data was collected to develop the CWQGPAqL were screened against the CCME (2007) data classification criteria. A summary of the collected data, as well as the qualifier for each study, are compiled in Appendix A. Many of the studies listed in Appendix A were utilized by the British Columbia Ministry of Water, Land and Air Protection (BC MOE) for the derivation of a water quality guideline for boron (1.2 mg/L) for the protection of aquatic life (Moss and Nagpal, 2003). Study classifications (primary, secondary, unacceptable) utilized by British Columbia (Moss and Nagpal, 2003) are those utilized by CCME (2007). For this reason, study classifications found in Moss and Nagpal (2003) were adopted if available or the original paper was reviewed to determine data quality. In general, primary toxicity studies involved acceptable test procedures, conditions and controls, measured toxicant concentrations and flow-through or renewal exposure conditions. Secondary toxicity studies usually involved unmeasured toxicant concentrations, static bioassay conditions, controls, pseudoreplication and may employ a wider array of methods. Unclassified toxicity information was also obtained from the BC MOE Ambient Water Quality Guidelines document for boron (Moss and Nagpal, 2003). For CWQGPAqL development, the original scientific publications were obtained for review and classification purposes, where possible. In the case

Scientific Criteria Document for the Development of a Canadian Water Quality Guideline for Boron

where it was not possible to retrieve the original publications, this unclassified toxicity information obtained from Moss and Nagpal (2003) was classified as secondary quality by default.

Acute toxicity studies generally involved test durations of 96 hours or less for vertebrates or 48 hours or less for invertebrates (the CCME 2007 protocol indicates that 24-96h exposures are short term exposures). Chronic toxicity data studies include complete life cycle tests and partial life cycle tests involving early life stages (the CCME 2007 protocol indicates that long-term exposure is defined as being ≥ 7 d for fish and invertebrates and ≥ 24 h for aquatic plants and algae).

2.1 Acute Toxicity

2.1.1 Vertebrates

Primary acute toxicity data for animals exposed to boron were available for early life stages of largemouth bass (*Micropterus salmoides*), fathead minnow (*Pimephales promelas*), rainbow trout (*Oncorhynchus mykiss*) and coho salmon (*Oncorhynchus kisutch*). Endpoints ranged from a 3 day NOEC (teratogenesis at hatching) of 0.109 mg/L for embryo-hatching stages of largemouth bass (Black *et al.*, 1993) to a 7 day EC₅₀ (embryo test) of 969 mg/L for rainbow trout (MELP, 1996). The reported effect concentrations for fathead minnow were a 48 and 96 hour LC₅₀ of 348 and 64.3 mg/L, respectively (MOE, 2007). The lowest reported 96 hour LC₅₀ values for coho salmon and rainbow trout were 304.1 mg/L (MELP, 1996) and 275 mg/L (MOE, 2007), respectively.

Secondary acute toxicity data were available for embryo and juvenile life stages of frog (*Xenopus laevis*), bluegill sunfish (*Lepomis macrochirus*), zebrafish (*Danio rerio*), bonytail (*Gila elegans*), Colorado squawfish (*Ptychocheilus lucius*), razorback sucker (*Xyrauchen texanus*), mosquito fish (*Gambusia affinis*), coho salmon (*Oncorhynchus kisutch*), and chinook salmon (*Oncorhynchus tshawytscha*). The lowest effect concentration was a 24 hour TL_m of 4.6 mg/L for the bluegill sunfish (Turnbull *et al.*, 1954), whereas the highest effect concentration was 24-hour TL_m of 3,145 mg/L for the mosquito fish (Wallen *et al.*, 1957). The following lists the lowest reported endpoints for the frog (4-d NOEC or 35.9 mg/L) (Fort *et al.*, 1999), and mosquito fish (96-h NOEC of <204 mg/L) (Wallen *et al.*, 1957). The lowest reported 96-hour LC₅₀ values for bonytail, Colorado squawfish and razorback sucker were 280, 279 and 233 mg/L, respectively (Hamilton, 1995). Hamilton and Buhl (1990) reported the lowest 96 hours LC₅₀ values for coho and Chinook salmon of 447 and 566 mg/L, respectively. Rowe *et al.* (1998) investigated 96 hour embryonic mortality of zygote zebrafish. A LOEC of 99.5 mg/L was reported based on 88% mortality (Rowe *et al.*, 1998).

Unclassified acute toxicity data were available for the rainbow trout, minnow and toad (*Bufo vulgaris*). The lowest LC₅₀ value for the rainbow trout was 339 mg/L (48 hours) (Sprague, 1972 in Birge and Black, 1977;). A minnow (species not identified) experienced a median lethal dose (TL_m or LC₅₀) at boron concentrations ranging from 340 to 374 mg/L with an unknown exposure duration (McKee and Wolf, 1963). The lowest effect concentration for toad embryos was 874 mg/L for malformations occurring after 24 hours of boric acid exposure (U.S. EPA, 1975).

2.1.2 Invertebrates

Primary acute toxicity data for invertebrates exposed to boron were available for the amphipod (*Hyalella azteca*), water flea (*Daphnia magna*), and midges (*Chironomus decorus*, *C. tentans*). 48 hour LC₅₀ concentrations ranged from 21.3 mg/L for neonate water fleas (MELP, 1996) to 1,376 mg/L for larvae midge *C. decorus* (Maier and Knight, 1991). The lowest 96 hour LC₅₀ for the amphipod was 28.9 mg/L (MELP, 1996).

Secondary acute toxicity data was available for the water flea, where neonates exposed to boron displayed a 48-h LC₅₀ of 133 mg/L (Gersich, 1984).

Kapu and Schaeffer (1991) examined the effects of boron exposure to the planarian (*Dugesia dorotocephala*) under static conditions with unmeasured concentrations. Effect concentrations ranged from 1 to 10 mg/L for hyperkinesias, spiraling and head/nose twist within five minutes of boron exposure. Toxicity data reported by Kapu and Schaeffer (1991) were classified as unacceptable as toxicity endpoints were evaluated using an arbitrary ranking scale for behavioural abnormalities and the responses observed at 1 to 5 minutes were not observed at 10 to 30 minutes or 40 to 60 minutes during the 1 hour behavioural test.

Unclassified acute toxicity data were available for the protozoan (*Entosiphon sulcatum*), mosquito larvae (*Anopheles quadrimaculatus*) and three unspecified species of mosquito larvae. Effect concentrations ranged from a threshold concentration of 1 mg/L for a reduction in cell replication for the protozoan (Bringmann, 1978) to a 48 hour LC₁₀₀ of 2,797 mg/L for pupae of three mosquito larvae (U.S. EPA, 1975). The lowest effect concentration for mosquito larvae was 92% mortality after a 48 hour exposure to a boron concentration of 25 mg/L (Fay, 1959). Gersich (1984) exposed water fleas to boric acid and reported a 48 hour LC₅₀ of 133 mg/L.

2.2 Chronic Toxicity

2.2.1 Vertebrates

Primary chronic toxicity data for vertebrates were available for rainbow trout, largemouth bass and fathead minnow. Black *et al.* (1993) exposed embryo and larval stage rainbow trout and largemouth bass to boric acid under flow through conditions with measured concentrations. Effect concentrations for embryo-larval rainbow trout ranged from a 32 day LOEC (mortality) of 0.1 mg/L to a 32 day LC₅₀ of 138 mg/L (Black *et al.*, 1993). The lowest effect concentration for embryo-larval stage largemouth bass was an 11 day LOEC (mortality) of 12.17 mg/L, while the highest was an 11 day LC₅₀ of 92 mg/L (Black *et al.*, 1993). Larval fathead minnow (*Pimephales promelas*) were exposed to boron for a 7-d period using a static renewal exposure, with a resulting IC₂₅ (growth inhibition) of 20.6 (8.5-26.5 mg/L 95% CI)(MOE 2007, unpublished data).

Secondary chronic toxicity data for vertebrates were available for embryo larval stages of rainbow trout, goldfish (*Carassius auratus*), channel catfish (*Ictalurus punctatus*), leopard frog (*Rana pipiens*), Fowler's toad (*Bufo fowleria*), as well as for blastula life-stage eggs of the wood frog (*Rana sylvatica*), Jefferson salamander (*Ambystoma jeffersonianum*), spotted salamander (*Ambystoma maculatum*), American toad (*Bufo americanus*) and alevin/fry coho salmon. The majority of the secondary toxicity data was reported by Birge and Black (1977) who exposed early life stages of various fish and amphibian species under continuous flow-through conditions to boric acid and borax at two different levels of water hardness (50 or 200 mg CaCO₃/L). The Scientific Criteria Document for the Development of a Canadian Water Quality Guideline for Boron

lowest effect concentration reported by Birge and Black (1977) was a 28 day LOEC of 0.01 mg/L for embryo-larval stages of rainbow trout. The lowest LOEC values for channel catfish (9 day), goldfish (7 day), leopard frog (7 day), and Fowler's toad (7 day) were 1, 8.33, 9.6, and 53.5 mg/L, respectively (Birge and Black, 1977). A study conducted by Davis and Mason (1973) which exposed alevins and fry coho salmon to sodium metaborate reported a 23 day LC₅₀ of 93 mg/L. Laposta and Duncan (1998) exposed various amphibian eggs to sodium tetraborate under static conditions with measured concentrations. The lowest effect concentration for wood frog, Jefferson salamander and spotted salamander blastula life-stage eggs was 50 mg/L for deformed hatchlings (Laposta and Duncan, 1998). Blastula life-stage American toad eggs also had a LOEC of 50 mg/L for the proportion of eggs hatching. Rowe *et al.* (1998) reported a 6 week LOEC of 108 mg/L for embryonic mortality of embryo rainbow trout exposed to boric acid.

Unclassified chronic toxicity data were available for early life stages of rainbow trout, largemouth bass, zebrafish (*Brachydanio rerio*), and the fathead minnow. The lowest effect concentration for the rainbow trout, largemouth bass, zebrafish and fathead minnow was a 32 day LOEC of 0.1 mg/L (Birge and Black, 1981), an 11 day LOEC of 12.17 mg/L (Birge and Black, 1981), a 32 day MATC of 10.04 mg/L (Hooftman *et al.*, 2000), and a 32 day LOEC for growth of 24 mg/L (Procter and Gamble, unpublished), respectively.

Of all of the species and life stages investigated in boron aquatic toxicity studies, the early life stages of rainbow trout appear to be the most sensitive in chronic exposures, based on the data presented in Appendix A. Considerable variability is observed between the no-effect and lowest effect concentrations of boron in studies conducted with embryo-larval stages of the trout (Birge and Black, 1977; Black *et al.*, 1993). Consistent concentration-related lowest observed effect levels in these studies for early life stages of the rainbow trout range from <0.1 to >17.0 mg/L (Figure 2). The flat dose-response curve observed for boron over the lower end of the exposure range may have complicated the determination of precise no-effect and lowest effect values (Black *et al.*, 1993). For instance, in studies conducted by Birge and Black (1977), it was common to see control-adjusted effects that ranged from two to eight percent for exposure concentrations spanning two to three orders of magnitude (Figure 1).

The flat dose-response curve may be partially attributed to the endpoints utilized. For instance, the frequency of teratogenesis included in studies conducted by Birge and Black (1977) have not been found to increase as proportionately with exposure concentrations as does mortality. Therefore, for trout stages exposed to boron it appears that the threshold effect for toxicity may be observed over a rather extensive concentration range (Black *et al.*, 1993). Given the variability of the test response data, only consistent dose-response effects (such as mortality) should be considered in the development of aquatic guidelines for boron (Black *et al.*, 1993).




Figure 1 The Effects of Boric Acid Administered to Embryo-Larval Stages of Trout in Soft Water (Birge and Black, 1977)

Compared with other research, the Birge and Black studies have consistently found very low concentration toxicity levels for a variety of aquatic species. However, other scientists and studies have not been able to reproduce these values using similar conditions and species (Moss and Nagpal, 2003). The toxicity data points of concern include the 28 day LOEC of 1.0 mg/L from Birge and Black (1977) and the 32 day LOEC of 0.1 mg B/L (Birge and Black, 1981) for early life-stage rainbow trout. Similar low effect concentrations for rainbow trout were observed in Black *et al.* (1993) when using reconstituted water (hardness = 188±10 mg/L as CaCO₃, alkalinity = 54±7 mg/L as CaCO₃, pH = 7.7±0.2), but not when using well water (hardness = 24-39 mg/L as CaCO₃, alkalinity = 25-38 mg/L as CaCO₃, pH = 6.8-7.1). In the case of exposures with well water, no significant adverse effects were observed following an 87-day exposure with embryo trout and larvae to measured boron concentrations ranging from 2.1 to 18 mg/L (Black *et al.* 1993). A further study by Birge *et al.* (1983) found that 5% of rainbow trout had boron induced teratogenesis at levels of 0.001 mg/L. Moss and Nagpal (2003) found that these specific data points for rainbow trout fell below the BC MOE guideline for boron for the protection of aquatic life, and were significantly lower than the other toxicity values for vertebrates. Therefore, Moss and Nagpal (2003) considered these data points to be outliers and they were not considered in the development of the British Columbia guideline. These outliers also were not considered for CWQGPAqL development.

2.2.2 Invertebrates

Primary chronic toxicity data were available for the water flea (*D. magna*) and midge (*C. decorus*). Effect concentrations ranged between a 21 day LOEC (mean brood size) of 13 mg/L to a 21 day LC₅₀ of 53.2 mg/L for the water flea (Lewis and Valentine, 1981). The lowest effect concentration for larval midge was a 96 hour LOEC (growth rate) of 20 mg/L (Maier and Knight, 1991).

Secondary chronic toxicity data was also available for the water flea *Daphnia magna* and *Ceriodaphnia dubia*, the amphipod (*Hyaella azteca*), and for a protozoa (*Opercularia bimarginata*). The lowest effect concentrations for *D. magna* and *C. dubia* was a 21 day MATC for growth of 4.67 mg/L (ANZECC, 2000) and a 14 day MATC for growth/reproduction of 13.4 mg/L (Hickey, 1989). Two 7 day LC₅₀s were reported for the amphipod, being 2.935 and >3.15

mg/L, respectively, for soft water and tap water (Borgmann *et al.*, 2005). A 72 hour NOEC of 10 mg/L was reported for the protozoa (Guhl, 1992b).

Unclassified chronic toxicity data for aquatic vertebrates were available for the water flea (*D. magna*) and for two protozoan species (*Entosiphon sulcatum* and *Paramecium caudatum*). The lowest effect concentration for the water flea was a threshold concentration for immobilization of <0.38 mg/L (McKee and Wolf, 1963).

2.2.3 Other Organisms (Algae, Macrophytes, etc.)

Primary chronic toxicity data were available for 4 to 7 day old green algae (*Selenastrum capricornutum*) (MELP, 1997). Algae were exposed to boron for 72 hour under static conditions. The lowest reported effect concentration was a 72 hour LOEC of 12.3 mg/L. Secondary toxicity data were available for the blue green algae (*Anacystis nidulans*), the green algae (*Chlorella pyrenoidosa*), the duckweed (*Spirodella polyrrhiza*), American waterweed (*Elodea canadensis*), stream water-crowfoot (*Ranunculus penicillatus*), water milfoil (*Myriophyllum alterniflorum*), common reed (*Phragmites australis*), the duckweed (*Lemna minor*), chlorella algae (*Chlorella vulgaris*), the green algae (*Scenedesmus subpicatus*), and the spiked or Eurasian watermillfoil (*Myriophyllum spicatum*). The effect concentrations range from a 10 day LOEC of 3.5 mg/L for frond production for the duckweed (*Spirodella polyrrhiza*) (Davis *et al.*, 2002) to a 32 day EC50 of 171 mg/L for the spiked or Eurasian milfoil (ANZECC, 2000).

Unclassified chronic toxicity data for algae and macrophytes were available. Endpoints for the green algae *C. pyrenoidosa* ranged from a growth or cell composition NOEC of 10 mg/L (Fernandez *et al.*, 1984) to a 72 hour IC of 100 mg/L (Maseo *et al.*, 1985). A 32 day IC50 of 40.3 mg/L for root growth was available for the spiked or Eurasian milfoil (Stanley, 1974).

2.3 Summary of Toxicity Data

The ranges of toxicity exhibited by aquatic organisms exposed to boron are summarized in Figure 2.

Primary acute toxicity data for vertebrates were available for early life stages of four fish species. Endpoints ranged from a 3 day NOEC (teratogenesis at hatching) of 0.109 mg/L for embryo-hatching stage largemouth bass (Black *et al.*, 1993) to a 7 day EC₅₀ (embryo test) of 969 mg/L for rainbow trout (MELP, 1996). Secondary and unclassified toxicity data were similar to the reported primary data with effect concentrations ranging between 4.6 and 3,145 mg/L.

Primary acute toxicity data for invertebrates were available for four species. The 48 hour LC₅₀ concentrations ranged from 21.3 mg/L for neonate water fleas (MELP, 1996) to 1,376 mg/L for larval midge (Maier and Knight, 1991). Secondary, unclassified and unacceptable toxicity data were similar to the reported primary data with effect concentrations ranging between 1 and 2,797 mg/L.

Primary chronic toxicity data for vertebrates were available for three fish species. Excluding the outliers identified in Section 2.2, effect concentrations ranged from a 36 day LOEC (mortality) of 1.34 mg/L for embryo-larval rainbow trout to a 32 day LC₅₀ of 138 mg/L for rainbow trout (Black *et al.*, 1993). Lower effect concentrations (0.01 mg/L) for embryo-larval rainbow trout were observed in the unclassified toxicity studies (Birge and Black, 1977).

Primary chronic toxicity data for invertebrates were available for two species. Effect concentrations ranged between a 21 day LOEC (mean brood size) of 13 mg/L to a 21 day LC₅₀ of 53.2 mg/L for the water flea (Lewis and Valentine, 1981). Effect concentrations for additional unclassified studies differed from those reported by the primary studies, with a range of <0.38 to ~266 mg/L.

Primary chronic toxicity data were available for 4 to 7 day old green algae (*Selenastrum capricornutum*) (MELP, 1997). The lowest reported effect concentration for algae exposed to boron was a 72 hour LOEC of 12.3 mg/L. Effect concentrations of secondary and unclassified toxicity data ranged from 1 to ~171 mg/L.

Additional factors such as water quality parameters and essentiality may also affect the toxicity of boron to aquatic species. These issues are discussed in Sections 2.4 and 2.5.

2.4 Effects of Water Quality Parameters on Toxicity

The toxicity of many elements depends on the chemical speciation of the dissolved species. Boron was added as several water soluble forms for aquatic toxicity studies (Appendix A) such as boric acid, sodium tetraborate, sodium borate, borax, sodium metaborate and sodium perborate. Birge and Black (1977) conducted a comparative toxicity test of boron administered as borax or boric acid. When the LC₁ and LC₅₀ data for all species were combined, there was no statistically significant difference between the toxicity values for boric acid and borax. However, this does not hold true for all water soluble forms of boron. For instance, Turnbull *et al.* (1954) examined the acute toxicity of sodium metaborate and boron trifluoride on bluegill sunfish. The 24 hour TLM (median tolerance limit) for sodium metaborate and boron trifluoride were 4.6 and 2,389 mg/L, respectively (Turnbull *et al.*, 1954). Furthermore, the 24 hour TLM for boric acid and sodium tetraborate were 3,145 and 1,360 mg/L, respectively, for adult female mosquito fish (*Gambusia affinis*). Although there were no statistically significant differences

between the aquatic toxicity of boric acid and borax, this may not hold true for other water soluble forms of boron.

The toxicity of metals to aquatic organisms is often modified by water hardness. In general, there does not appear to be any significant interaction between water hardness and boron toxicity (Birge and Black, 1977; Laws, 1981; Hamilton and Buhl, 1990; Maier and Knight, 1991). Birge and Black (1977) investigated the effect of water hardness (at 50 and 200 mg CaCO₃/L) on the toxicity of boron to aquatic organisms. When LC₁ and LC₅₀ data for all species were combined, there were no statistically significant differences between the values for boric acid and borax. The hardness of the test medium was found to exert an effect on the toxicity of boron to fish; however, this effect was not consistent. For instance, borax exhibited greater toxicity in hard water to embryos and larvae of both goldfish and channel catfish, while embryos and larvae of rainbow trout were more sensitive to borax in soft water. Maier and Knight (1991) examined the interaction of boron and water hardness on the mortality of neonate *Daphnia magna*. No significant interaction was observed between boron toxicity and water hardness, as the mortality (48 hour LC₅₀) associated with boron exposure did not significantly change when water hardness increased from 10.6 to 170 mg/L. British Columbia MELP (1996) found no correlation between hardness and toxicity for coho salmon, rainbow trout and chironomid test outcomes. However, decreasing toxicity with increased hardness was observed for *Daphnia* and *Hyaella* (MELP, 1996). Overall, water hardness does not significantly impact the toxicity of boron to aquatic organisms.

Laws (1981) found that there was no interaction between sulphate and boron in natural aquatic ecosystems.

Several studies have shown that the low-level effects observed in reconstituted laboratory water are not predictive of the much higher effect levels found under natural water exposure conditions (Butterwick *et al.*, 1989; Black *et al.*, 1993). Consistent concentration-related lowest observed effect levels in reconstituted laboratory water range from <0.1 to greater than 18.0 mg/L for rainbow trout (Birge and Black, 1977). The lowest effect concentration for rainbow trout exposed to natural water amended with boron from the Erwin hatchery, Brookville Lake and the Firehole River was 1.0 mg/L; however, ambient concentrations of 0.75 mg/L did not produce significant effects (Black *et al.*, 1993). No effects were observed at measured boron concentrations of 2.1 mg/L administered for 87 days during early life stage trout studies. In addition, a boron concentration of 18 mg/L produced no significant effects when exposure occurred 20 days after egg fertilization and maintained until 60 days post-hatching (Black *et al.*, 1993). Boron concentrations were measured in California streams where viable populations of wild trout were observed (Black *et al.*, 1993). In approximately 88% of the streams, boron was detected at or below 0.5 mg/L. However, in one stream, boron was found at concentrations as high as 13 mg/L (Black *et al.*, 1993). These field observations further support that rainbow trout are less sensitive to natural water boron concentrations than when exposed to boron in laboratory reconstituted water. These variations may be a result of differences in the natural chemical composition of waters from different geographical regions which may modify the toxicity of boron (Black *et al.*, 1993). However, the specific component of the water chemistry responsible for toxicity modification is unknown. It is hypothesized that boron attenuation may occur *via* complexation with organic compounds or adsorption to particulate matter.

2.5 Essentiality and Deficiency

The essentiality of chemicals should be considered during water quality guideline derivation for the protection of aquatic life. Essentiality of an element means that the absence or deficiency of the element results in the impairment of life functions, and that the impairment can be prevented or corrected only by supplementation of physiological levels of the element and not by others. Essential chemicals are different than non-essential chemicals as negative effects on organisms are observed when insufficient levels (*i.e.*, levels below the compensation limit of accumulation / assimilation of the organism) of the chemical are present in the environment. This deficiency varies between organisms, between species, and within species based on respective locale (adaptation). As organisms have adapted to their natural habitat, it can be assumed that the natural background concentrations of essential chemicals at a given locale fulfill the requirements of essentiality to organisms there. Organisms requiring levels of essential chemicals in greater quantities than those naturally present in a particular environment (natural background concentrations) are not expected to be present naturally in this environment to begin with, or if present, would suffer from deficiency not caused by anthropogenic influences.

The effects of boron have been studied on a variety of freshwater fish, amphibians, invertebrates and plants. At lower concentrations, boron has been found to be beneficial to some freshwater organisms. For instance, the addition of 0.4 mg B/L to ponds used for raising carp increased production by 7.6% (Avetisyan, 1983). Furthermore, Fort *et al.* (1999) found that boron was nutritionally essential for the reproduction and development in frogs (*Xenopus laevis*).

Rowe *et al.* (1998) found that the shape of the dose-response curve in rainbow trout (*Oncorhynchus mykiss*) and zebrafish follows the U-shaped adverse response of an essential nutrient. This shape reflects effects of exposure to boron concentrations below the level to meet physiological requirements and toxic effects due to exposure to high concentrations of boron that exceed the threshold for safety. For rainbow trout embryos, chronic exposures below 9.7×10^{-2} mg/L impaired embryonic growth. In addition, zebrafish exposed to boron concentrations below 2.2×10^{-3} mg/L experienced zygote death (Rowe *et al.*, 1998). The theory regarding boron essentiality suggested by the results of Rowe *et al.* (1998) may explain why the Birge and Black studies have consistently found very low concentration toxicity levels for boron for a variety of aquatic species.

In order to prevent adverse health effects to organisms caused by a deficiency of essential chemicals, recommended threshold levels for these chemicals should not fall below the level required by the organism to remain healthy.

3.0 BIOACCUMULATION

The bioaccumulation and biomagnification of boron in the aquatic environment is not clearly understood (Moss and Nagpal, 2003). In general, the literature suggests that aquatic environments are not likely to experience boron biomagnifications, however, there is evidence of bioaccumulation (Wren *et al.*, 1983; Butterwick *et al.*, 1989). Boron has been shown to accumulate in aquatic plants (Wren *et al.*, 1983; Saiki *et al.*, 1993), although this is species specific. Fernandez *et al.* (1984) examined the bioaccumulation of boron in green alga (*Chlorella pyrenoidosa*). After seven days, the bioconcentration factor (BCF) for three concentration levels ranged from four to five. In a study conducted by Davis *et al.* (2002), duckweed (*Spirodella polyrrhiza*) did not remove significant amounts of boron from the

treatment solutions. However, Glandon and McNabb (1978) found that duckweed (*Lemna minor*) did bioaccumulate boron compared to other hydrophytes (e.g., *Ceratophyllum demersum*). A curvilinear relationship was observed between ambient boron concentrations and concentrations in the plant tissues, suggesting that both active and passive transport of boron across plant root membranes occurs in this species. Frick (1985) found that pH affected the bioaccumulation of boron by duckweed (*Lemna minor*).

Despite a tendency to accumulate in plants and algae, boron bioaccumulation by invertebrates and fish has been shown to be lower (Saiki et al 1993), and boron does not appear to biomagnify through the food chain (Wren et al., 1983; Saiki et al., 1993). The BCFs for boron in marine and freshwater plants, fish and invertebrates were estimated to be less than 100 (Thompson et al., 1972). Experimental BCFs for fish have ranged between 52 and 198 (Tsui and McCart, 1981). In the marine environment, Thompson et al. (1976) found no evidence of boron bioaccumulation in sockeye salmon (*Oncorhynchus nerka*) tissues or pacific oyster (*Crassostrea gigas*). In both instances, tissue boron concentrations approximated boron concentrations in the test water (Thompson et al., 1976). In the oyster, tissue concentrations returned to background levels by the 71st day of the study after boron exposure ceased, indicating a fairly rapid clearance of boron. Furthermore, a study conducted by Suloway et al. (1983) examined the bioaccumulation potential of the components of coal fly ash extract in fathead minnows (*Pimephales promelas*) and green sunfish (*Lepomis cyanellus*). The BCF for boron was 0.3 for both species. These BCF values suggest that boron does not significantly bioconcentrate or biomagnify in the aquatic environment.

4.0 IMPACT ON TASTE AND ODOUR OF WATER AND FISH TAINING

No information on the impact of boron on the taste and odour of water or fish tainting was available. Neither Health Canada (1990) nor WHO (2003) derived a drinking water quality guideline for boron based on this endpoint.

5.0 MUTAGENICITY

Boron and its compounds are not considered mutagenic in either bacterial or mammalian systems.

In a *Salmonella typhimurium*-mammalian microsome mutagenicity assay, boron did not enhance or inhibit the activity of benzo[a]pyrene, a known mutagen (Benson et al., 1984). Boric acid and borax were not mutagenic in *Salmonella typhimurium* with or without rat or hamster S9 fraction (Haworth et al., 1983; Benson et al., 1984; NTP, 1987; Steward, 1991). Boric acid was not mutagenic, or produced equivocal results, in the streptomycin-dependant *Escherichia coli* Sd-4 assay (Demerec et al., 1951; Iyer and Szybalski, 1958; Szybalski, 1958). Sodium perborate was shown to interact with DNA in an *Escherichia coli* Pol A assay, presumably by being converted to hydrogen peroxide (Rosenkranz, 1973). A single study (Odunola, 1997) found that boric acid induced *B*-galactosidase synthesis (a response to DNA lesions) in *E. coli* PQ37 (SOS chromotest) with and without metabolic activation.

Results in mammalian systems for mutagenicity were all negative (U.S. EPA, 2004). Boric acid did not induce unscheduled DNA synthesis in primary cultures of male rat hepatocytes (Bakke, 1991). Refined borax, crude borax ore and kermite ore were not mutagenic in V79 Chinese hamster cells, mouse embryo fibroblasts or diploid human foreskin fibroblasts (Landolph, 1985). Scientific Criteria Document for the Development of a Canadian Water Quality Guideline for Boron 21

In addition, other tests have shown that boric acid does not induce chromosomal aberrations or sister chromatid exchanges in Chinese hamster ovary cells (NTP, 1987). Forward mutations in mouse lymphoma cells were not induced by boric acid with or without rat liver S9 (NTP, 1987; McGregor *et al.*, 1988). In addition boric acid did not induce mutations at the thymidine kinase locus in mouse lymphoma cells in the presence or absence of rat liver activation system (Rudd, 1991). Finally, boric acid did not induce chromosomal or mitotic spindle abnormalities in bone marrow erythrocytes in the micronucleus assay in Swiss-Webster mice (O'Loughlin, 1991).

6.0 DERMAL AND OTHER EFFECTS

Absorption of boron is poor through intact skin, but is much greater through damaged skin (WHO, 2003). Little information was found on the protection of recreational water uses based on public health concerns, wildlife protection, toxicant interactions or sediment quality. Australia and New Zealand Environment and Conservation Council (ANZECC, 2000) have a guideline for protection of recreational water uses and aesthetics of 1 mg/L, although the basis of this guideline was not provided.

7.0 METHOD DETECTION LIMITS

The Ontario Ministry of the Environment uses a method to measure trace metals (boron) in water that has been certified by the Canadian Association for Environmental Analytical Laboratories (CAEAL). The method (E3474) determines the concentration of trace metals in surface and groundwater by dynamic reaction cell (DRC) using inductively coupled plasma-mass spectrometry (ICP-MS) (R. Moody, Manager Spectroscopy Section, Laboratory Services Branch, Ontario Ministry of the Environment, personal communication). The current MOE laboratory detection limit in water is 0.2 $\mu\text{g L}^{-1}$ or 0.0002 mg L^{-1} . Environment Canada's National Laboratory for Environmental Testing (NLET) analyzes for trace metals (boron) in water using ICP-MS with a method detection limit of 0.5 $\mu\text{g L}^{-1}$ or 0.0005 mg L^{-1} (J. Carrier, Inorganics Analyst, NLET, Environment Canada, personal communication). Various analytical methods are available to accurately detect boron levels in water ranging from 0.01 mg L^{-1} to 10 mg L^{-1} (Health Canada, 1990).

8.0 DERIVATION OF SPECIES SENSITIVITY DISTRIBUTIONS

Species sensitivity distributions (SSDs) represent the variation in sensitivity of species to a substance by a statistical or empirical distribution function of responses for a sample of species. The basic assumption of the SSD concept is that the sensitivities of a set of species can be described by some distribution, usually a parametric sigmoidal cumulative distribution function. The use of SSDs has become common in ecological risk assessment as well as environmental quality guideline development. In 2007, the Canadian Council of Ministers of the Environment (CCME) established a new protocol for deriving water quality guidelines for the protection of aquatic life. Canadian Water Quality Guidelines for the Protection of Aquatic Life (CWQGs-PAL) are nationally accepted threshold values for substances and other attributes (such as pH and temperature) in water. These values are determined such that no adverse toxic effects are expected in aquatic plants and animals. The guidelines are management tools constructed to ensure that anthropogenic stresses, such as the introduction of toxic substances, do not result in the degradation of Canadian waters. The 2007 protocol for the derivation of CWQGs-PAL now provides three options for guideline development, all dependent on the quantity and quality of toxicity data available. The first, and recommended, guideline derivation method involves

modeling the cumulative species sensitivity distribution (SSD) with estimating the 95% confidence interval. The guideline is defined as the intercept of the 5th percentile of the species sensitivity distribution (CCME, 2007) and is interpreted as protecting 95% of species. This level of protection ensures that aquatic community structure and function is maintained. SSD derived guidelines are referred to as Type A guidelines. If data are inadequate to derive a Type A guideline, then either a Type B1 or B2 guideline can be set, where an assessment or uncertainty factor is applied to the lowest effect concentration found in the scientific literature. The minimum data requirements for setting either a Type A, B1 or B2 guideline can be found in the CCME (2007) protocol document. The Type B1 and B2 guidelines are a modified version of the method traditionally used to derive Canadian Water Quality Guidelines for the Protection of Aquatic Life described in the 1991 CCME protocol. In the case of boron, suitable short-term and long-term datasets were provided for the development of a Type A guideline. Freshwater SSDs for freshwater biota were derived for both exposure durations following the CCME Protocol for the Derivation of Water Quality Guidelines for the Protection of Aquatic Life (CCME, 2007). Results may be used in accordance with the protocol to derive guideline values for boron.

To generate the short-term and long-term SSDs primary, secondary and unclassified toxicity data were included; however, datapoints classified as unacceptable were excluded. When multiple data points or effects (*e.g.*, growth, mortality, reproduction) were available for the same species professional judgment was utilized to select a representative species effect concentration (*e.g.*, lowest value or geometric mean). Using a customized Microsoft Excel-based software package, SSD Master Version 2.0, a total of five cumulative distribution functions were fit to the data using regression techniques. Model fit was assessed using statistical and graphical techniques. The best model was selected based on goodness-of-fit and model feasibility. Model assumptions were verified graphically. The concentration of boron in freshwater at which 5% of species are predicted to be affected was determined for both short-term and long-term scenarios with 95% confidence intervals on the mean (expected) value.

The Generalized Linear Model (GLiM) framework described by Kerr and Meador (1996) and Bailer and Oris (1997) was used as the modelling framework to derive the short- and long-term SSDs. The framework involves using link functions to transform metrics and assigning appropriate error distributions (*e.g.*, binomial distribution for quantal responses). Each species for which appropriate toxicity data were available was ranked according to sensitivity (from lowest to highest value), and its centralized position on the SSD (Hazen plotting position) was determined using the following standard equation (Aldenberg *et al.*, 2002; Newman *et al.*, 2002):

$$\text{Hazen Plotting Position} = \frac{i - 0.5}{N}$$

where:

- i = the species rank based on ascending toxicity values
- N = the total number of species included in the SSD derivation

8.1 Short-term SSD Derivation

Short-term exposure benchmark concentrations are derived using severe effects data (such as lethality or immobilization) of defined short-term exposure periods (24-96h). These guidelines identify estimators of severe effects to the aquatic ecosystem and are intended to give guidance

on the impacts of severe, but transient, situations (e.g., spill events to aquatic receiving environments and infrequent releases of short-lived/nonpersistent substances). Short-term benchmark concentrations *do not* provide guidance on protective levels of a substance in the aquatic environment, as short-term benchmark concentrations are levels which *do not* protect against adverse effects.

The minimum data requirements for the development of a short-term Type A (SSD-derived) benchmark were met, and these are listed in Table 8-1.

Table 8- 1 Minimum Data Set Requirements for the Gereneration of a Short-Term Freshwater Benchmark Concentration

Toxicity tests required for the generation of a short-term SSD, broken out as follows:	
Fish:	3 tests on 3 different species including 1 salmonid, 1 non-salmonid.
Invertebrates:	3 tests on 3 different species including 1 planktonic crustacean, 2 others. For semi-aquatic invertebrates, the life stages tested must be aquatic. It is desirable, but not necessary, that one of the aquatic invertebrate species be either a mayfly, caddisfly, or stonefly.
Plant/Algae:	None (for non-phytotoxic substances), 2 (for phytotoxic substances).
Toxicity data for amphibians are highly desirable, but not necessary. Data must represent fully aquatic stages.	
Acceptable endpoints for acute guidance: LC/EC50 (severe effects) Note: Primary or secondary data are acceptable.	

A total of 13 data points (all LC₅₀ values) were used in the derivation of the benchmark concentration (Table 8-2). These 13 data points were retrieved from toxicity studies meeting the requirements for primary and secondary data, according to CCME (2007) protocol. Intra-species variability was accounted for by taking the geometric mean of the studies considered to represent the most sensitive lifestage and endpoint. Each data point was ranked according to sensitivity, and its centralized distribution on the species sensitivity distribution (SSD) was determined using the Hazen plotting position (estimate of the cumulative probability of a data point). The plotting positions are treated as observed proportions of species affected. These positional rankings, along with their corresponding LC₅₀ values, were used to derive the SSDs.

Table 8- 2 Short-term LC₅₀s for species exposed to boron in freshwater

<i>Rank</i>	<i>Scientific Name</i>	<i>Common Name</i>	<i>Endpoint</i>	<i>LC₅₀ (mg B/L)</i>	<i>Data Quality</i>	<i>Hazen Plotting Position</i>	<i>Reference</i>
1	<i>Lepomis macrochirus</i>	Bluegill	Mortality	4.6	S	0.04	Turnbull et al., 1954
2	<i>Pimephales promelas</i>	Fathead minnow	Mortality	64.3	P	0.12	MOE, 2007
3	<i>Daphnia magna</i>	Water flea	Mortality	101.2	Grouped P*, P, S	0.19	Maier & Knight, 1991; MOE 2007; MELP 1996; Lewis & Valentine 1981; Gersich, 1984
4	<i>Chironomus</i>	Midge	Mortality	136.7	Grouped	0.27	MELP, 1996

Rank	Scientific Name	Common Name	Endpoint	LC ₅₀ (mg B/L)	Data Quality	Hazen Plotting Position	Reference
	<i>tentans</i>				P*		
5	<i>Hyaella azteca</i>	Freshwater shrimp	Mortality	141.1	Grouped P*	0.35	MELP, 1996
6	<i>Xyrauchen texanus</i>	Razorback sucker	Mortality	233	S	0.42	Hamilton, 1995
7	<i>Ptychocheilus lucius</i>	Colorado squawfish	Mortality	279	S	0.50	Hamilton, 1995
8	<i>Gila elegans</i>	Bonytail	Mortality	280	S	0.58	Hamilton, 1995
9	<i>Oncorhynchus mykiss</i>	Rainbow trout	Mortality	351.7	Grouped P*, P	0.65	MELP, 1996; MOE 2007
10	<i>Oncorhynchus kisutch</i>	Chinook salmon	Mortality	372.9	Grouped P*	0.73	MELP, 1996
11	<i>Oncorhynchus tshawytscha</i>	Coho salmon	Mortality	566	S	0.81	Hamilton and Buhl, 1990
12	<i>Gambusia affinis</i>	Mosquitofish	Mortality	632	S	0.88	Wallen et al., 1957
13	<i>Chironomus decorus</i>	Midge	Mortality	1376	P	0.96	Maier and Knight, 1991

Data Quality:

S = Secondary; P = Primary; P* = Primary as defined by Moss and Nagpal (2003); S* = Secondary as defined by Moss and Nagpal (2003)

Grouped Indicates that the geomean of multiple values was used to calculate the effect concentration

The values reported in Table 8-2 range from a 24h-LC₅₀ of 4.6 mg/L for the Bluegill sunfish, *Lepomis macrochirus* (Turnbull et al., 1954), to a 48-h LC₅₀ of 1376 mg/L for the midge, *Chironomus decorus* (Maier and Knight, 1991). Geometric mean values were calculated for *Daphnia magna*, *Chironomus tentans*, *Hyaella azteca*, *Oncorhynchus mykiss*, and *Oncorhynchus kisutch* (Table 8-3). Effect concentrations reported for the remaining species were taken from single studies.

Table 8- 3 Studies Used to Derive Geometric Means for Short-Term Data

Organism	Endpoint	Effect Concentration (mg/L)	Geometric Mean (mg/L)	Reference
<i>Daphnia magna</i> (Water flea)	48-h LC ₅₀	141 165 21.3 52.4 139.2 226 133	101.2	Maier & Knight, 1991; MOE 2007; MELP 1996; Lewis & Valentine 1981; Gersich, 1984
<i>Chironomus tentans</i> (Midge)	96-h LC ₅₀	118 137.7 157.3	136.7	MELP, 1996
<i>Hyaella azteca</i> (Freshwater shrimp)	96-h LC ₅₀	28.9 291.3 333.6	141.1	MELP, 1996
<i>Oncorhynchus</i>	96-h LC ₅₀	275	351.7	MELP, 1996;

<i>mykiss</i> (Rainbow trout)		336 379.6 436.2		MOE 2007
<i>Oncorhynchus kisutch</i> (Chinook salmon)	96-h LC ₅₀	304.1 357.4 477.1	372.9	MELP, 1996

Five cumulative distribution functions (normal, logistic, Gompertz, Weibull, and Fisher-Tippett) were fit to the data, both in arithmetic space (no transformation of LC₅₀ values) and log space (log transformed LC₅₀ values) using regression methods. Model fit was assessed using statistical and graphical techniques. The best model was selected based on consideration of goodness-of-fit test and model feasibility. Model assumptions were verified graphically and with the use of statistical tests.

Of the ten models tested, the log-Gompertz model fit the data best (Figure 3). The Anderson-Darling Goodness of Fit test statistic (A^2) was 0.304 (P-value > 0.10). The equation of the fitted log-Gompertz model is of the form:

$$y = 1 - e^{-e^{\left(\frac{x-\mu}{\delta}\right)}}$$

Where x is the log (concentration) and y is the proportion of species affected. For the fitted model, $\mu = 2.5319$ and $\delta = 0.3609$. Summary statistics for the short-term SSD are presented in Table 8-4. The 5th percentile on the short-term SSD is 29 mg/L. The lower fiducial limit (5%) on the 5th percentile is 15 mg/L, and the upper fiducial limit (95%) on the 5th percentile is 55 mg/L. The benchmark concentration is defined as the 5th percentile on the SSD.

Therefore, the short-term exposure benchmark concentration indicating the potential for severe effects (e.g. lethality or immobilization) to sensitive freshwater life/marine life during transient events is 29 mg/L.

Table 8- 4 Short-term Concentration for Boron Resulting from the SSD Method

	Concentration
SSD 5th percentile	29 mg/L
Lower 95% confidence limit	15 mg/L
Upper 95% confidence limit	55 mg/L




Figure 3 SSD for boron in freshwater derived by fitting the log-Gompertz model to the logarithm of acceptable short-term LC₅₀s of thirteen aquatic species versus Hazen plotting position (proportion of species affected).

8.2 Long-term SSD Derivation

Long-term exposure guidelines identify benchmarks in the aquatic ecosystem that are intended to protect all forms of aquatic life for indefinite exposure periods (≥ 7 d exposures for fish and invertebrates, ≥ 24 h exposures for aquatic plants and algae).

The minimum data requirements for the development of a long-term Type A (SSD-derived) guideline were met, and these are listed in Table 8-5.

Table 8- 5 Minimum Data Set Requirements for the Generation of a long-term freshwater CWQG

Toxicity tests required for the generation of a long-term SSD, broken out as follows:
Fish: 3 tests on 3 different species including 1 salmonid, 1 non-salmonid.
Invertebrates: 3 tests on 3 different species including 1 planktonic crustacean, 2 others. For semi-aquatic invertebrates, the life stages tested must be aquatic. It is desirable, but not necessary, that one

of the aquatic invertebrate species be either a mayfly, caddisfly, or stonefly.
Plant/Algae: At least one study on a freshwater vascular plant or freshwater algal species (for non-phytotoxic substances), 3 studies (for phytotoxic substances)
Toxicity data for amphibians are highly desirable, but not necessary. Data must represent fully aquatic stages.
Acceptable endpoints for chronic guidance: Most appropriate ECx/ICx representing a no-effects threshold > EC10/IC10 > EC11-25/IC11-25 > MATC > NOEC > LOEC > EC26-49/IC26-49 > nonlethal EC50/IC50. Note: Primary or secondary data are acceptable.

A total of 28 data points (NOEC, EC₁₀, MATC and LOEC data) were used in the derivation of the long-term guideline (Table 4-2). Toxicity studies meeting the requirements for primary and secondary data, according to CCME (2007) protocol. Intra-species variability was accounted for by taking the geometric mean of the studies considered to represent the most sensitive lifestage and endpoint. Each data point was ranked according to sensitivity, and its centralized distribution on the species sensitivity distribution (SSD) was determined using the Hazen plotting position (estimate of the cumulative probability of a data point). The plotting positions are treated as observed proportions of species affected. These positional rankings, along with their corresponding no-effects and low-effects values, were used to derive the SSDs.

Insufficient EC₁₀ (one data-point) and EC₁₀₋₂₅ (no data) no-effect toxicity data were available to construct the SSD, therefore MATC and NOEC endpoints were predominantly used (in addition to the single EC₁₀ data-point). With respect to low effect data, insufficient EC₂₀ or EC₂₀₋₂₅ low-effect toxicity data were available, therefore LOEC endpoints were used.

Table 8- 6 Long-term low effects and no effects concentrations for species exposed to boron in freshwater

Rank	Scientific Name	Common Name	Endpoint	Effective Concentration (mg B/L)	Data Quality	Hazen Plotting Position	Reference
1	<i>Elodea canadensis</i>	American Waterweed	(NOEC)	1.0	S	0.01	ANZECC, 2000
2	<i>Spirodella polyrrhiza</i>	Duckweed	Fronnd production (10 d, MATC)	1.8	Grouped S	0.04	Davis et al., 2002
3	<i>Chlorella pyrenoidosa</i>	Green algae	Growth or cell composition (NOEC)	2.0	Grouped S	0.07	ANZECC, 2000; Fernandez et al., 1984
4	<i>Oncorhynchus mykiss</i>	Rainbow trout	Embryo survival at 60-d post-hatch (87 d, NOEC)	2.1	P	0.13	Black et al., 1993
5	<i>Ictalurus punctatus</i>	Channel catfish	(9 d, MATC)	2.4	Grouped S	0.16	Birge and Black, 1977
6	<i>Phragmites australis</i>	Common reed	Growth (4-month, NOEC)	4.0	S	0.19	Bergmann 1995
7	<i>Micropterus salmoides</i>	Largemouth bass	(11 d, MATC)	4.1	Grouped P	0.21	Black et al., 1993
8	<i>Chlorella vulgaris</i>	Green algae	Population growth (NOEC)	5.2	S	0.24	ANZECC, 2000
9	<i>Daphnia magna</i>	Water flea	Reproduction (NOEC)	6.0	S	0.27	ANZECC, 2000
10	<i>Opercularia bimarginata</i>	Unknown (Protozoa?)	Growth, reproduction (72-h NOEC)	10.0	S	0.33	Guhl 1992 (German), In Dyer 2001
11	<i>Brachydanio rerio</i>	Zebrafish	Mortality, growth, condition (34-d MATC)	10.0	S	0.36	Hoofman et al, 2000
12	<i>Selenastrum capricornutum</i>	Green algae	LOEC (72 hr)	12.3	P*	0.41	MELP, 1997 In Moss and Nagpal, 2003
13	<i>Ceriodaphnia dubia</i>	Water flea	Growth, reproduction (14-d MATC)	13.4	S	0.44	Hickey, 1989
14	<i>Entosiphon sulcatum</i>	Zooplankton	Growth (72-h NOEC)	15.0	S	0.47	Guhl 1992 (German), In Dyer 2001
15	<i>Carassius auratus</i>	Goldfish	(7 d, MATC)	15.6	Grouped S	0.50	Birge and Black, 1977
16	<i>Pimephales promelas</i>	Fathead minnow	Growth reduction (30 d, MATC)	18.3	Grouped S	0.53	Proctor and Gamble (unpublished) In Butterwick <i>et al.</i> , 1989 In Moss and Nagpal, 2003
17	<i>Chironomus decorus</i>	Midge	Growth (96-h NOEC)	20.0	S	0.59	Maier and Knight, 1991 In Dyer, 2001
18	<i>Paramecium caudatum</i>	Ciliate	Growth, reproduction (72-h NOEC)	20.0	S	0.61	Guhl 1992 (German), In Dyer 2001
19	<i>Rana pipiens</i>	Leopard frog	(7 d MATC)	20.4	Grouped S	0.56	Birge and Black, 1977 In

							Butterwick et al., 1989 and Eisler, 1990 In Moss and Nagpal, 2003
20	<i>Scenedesmus subspicatus</i>	Green algae	Growth (96-h EC 10)	30.0	S	0.70	Guhl 1992 (German), In Dyer 2001
21	<i>Myriophyllum spicatum</i>	Spiked or Eurasian water milfoil	(NOEC)	34.2	S	0.73	ANZECC, 2000
22	<i>Bufo fowleri</i>	Fowler's toad	(7 d, MATC)	48.6	S	0.79	Birge and Black, 1977
23	<i>Anacystis nidulans</i>	Blue-green algae	Growth or organic constituents (NOEC)	50.0	S	0.81	Martinez et al., 1986
24	<i>Bufo americanus</i>	American toad	proportion of eggs hatching (15-23 d, LOEC, 43% decrease compared to controls)	50.0	S	0.84	Laposata and Dunson, 1998
25	<i>Lemna minor</i>	Duckweed	Growth (7-d NOEC)	60.0	S	0.87	Wang, 1986
26	<i>Ambystoma jeffersonianum</i>	Jefferson's salamander	Proportion of deformed hatchlings (17-25 d, MATC increase in deformities compared to controls)	70.7	Grouped S	0.90	Laposata and Dunson, 1998
27	<i>Ambystoma maculatum</i>	Spotted salamander	Proportion of deformed Hatchlings (38-44 d, MATC, increase in deformities compared to controls)	70.7	Grouped S	0.93	Laposata and Dunson, 1998
28	<i>Rana sylvatica</i>	Wood frog	Proportion of eggs hatched (13-23 d, MATC)	70.7	Grouped S	0.99	Laposata and Dunson, 1998

Data Quality:

S = Secondary; P = Primary; P* = Primary as defined by Moss and Nagpal (2003); S* = Secondary as defined by Moss and Nagpal (2003); ? = Unclassified toxicity data classified as secondary quality by default (proprietary information, original publication could not be obtained)

Grouped Indicates that the geomean of multiple values was used to calculate the effect concentration

The values reported in Table 8-6 range from a NOEC of 1.0 mg/L for the American waterweed, *Elodea canadensis* (ANZECC, 2000), to a 13-22 d MATC of 70.7 mg/L for the Wood frog, *Rana sylvatica* (Laposata and Dunson, 1998). Geometric mean values were calculated for *Spirodella polyrrhiza*, *Clorella pyrenoidosa*, *Ictalurus punctatus*, *Micropterus salmoides*, *Carassius auratus*, *Pimephales promelas*, *Rana pipiens*, *Ambystoma jeffersonianum*, *Ambystoma maculatum*, and *Rana sylvatica* (Table 8-7). Effect concentrations reported for the remaining species were taken from single studies.

Table 8- 7 Studies Used to Derive Geometric Means for Long-Term Data

Organism	Endpoint	Effect Concentration (mg/L)	Geometric Mean (mg/L)	Reference
<i>Spirodella polyrrhiza</i>	10-d NOEC 10-d LOEC	0.9 3.5	1.8	Davis et al., 2002

Organism	Endpoint	Effect Concentration (mg/L)	Geometric Mean (mg/L)	Reference
(Duckweed)				
<i>Chlorella pyrenoidosa</i> (Green algae)	14-d NOEC 14-d NOEC	0.4 10	2.0	ANZECC, 2000; Fernandez et al., 1984
<i>Ictalurus punctatus</i> (Channel catfish)	9-d NOEC 9-d NOEC 9-d NOEC 9-d NOEC 9-d LOEC 9-d LOEC 9-d LOEC 9-d LOEC	0.49 0.75 1.01 9 1 1.04 5.42 25.9	2.4	Birge and Black, 1977
<i>Micropterus salmoides</i> (Largemouth bass)	11-d NOEC 11-d LOEC	1.39 12.17	4.1	Black et al., 1993
<i>Carassius auratus</i> (Goldfish)	7-d NOEC 7-d NOEC 7-d NOEC 7-d NOEC 7-d LOEC 7-d LOEC 7-d LOEC 7-d LOEC	6.8 8.53 9.2 26.5 8.33 22.5 27.33 48.75	15.6	Birge and Black, 1977
<i>Pimephales promelas</i> (Fathead minnow)	30-d NOEC 30-d LOEC	14 24	18.3	Proctor and Gamble (unpublished) In Butterwick <i>et al.</i> , 1989 In Moss and Nagpal, 2003
<i>Rana pipiens</i> (Leopard frog)	7-d NOEC 7-d NOEC 7-d NOEC 7-d NOEC 7-d LOEC 7-d LOEC 7-d LOEC 7-d LOEC	7.04 7.04 32.5 45.7 9.6 10.5 47.5 86	20.4	Birge and Black, 1977 In Butterwick et al., 1989 and Eisler, 1990 In Moss and Nagpal, 2003
<i>Ambystoma jeffersonianum</i> (Jefferson's Salamander)	17-25 d NOEC 17-25d LOEC	50 100	70.7	Laposata and Dunson, 1998
<i>Ambystoma maculatum</i> (Spotted salamander)	38-44 d NOEC 38-44 d LOEC	50 100	70.7	Laposata and Dunson, 1998

Organism	Endpoint	Effect Concentration (mg/L)	Geometric Mean (mg/L)	Reference
<i>Rana sylvatica</i> (Wood frog)	13-23 d NOEC 13-23 d LOEC	50 100	70.7	Laposata and Dunson, 1998

Five cumulative distribution functions (normal, logistic, Gompertz, Weibull, and Fisher-Tippett) were fit to the data, both in arithmetic space (no transformation of LC₅₀ values) and log space (log transformed LC₅₀ values) using regression methods. Model fit was assessed using statistical and graphical techniques. The best model was selected based on consideration of goodness-of-fit test and model feasibility. Model assumptions were verified graphically and with the use of statistical tests.

Of the ten models tested, the log-Normal distribution function (with a mean [μ] of 1.1443 and a standard deviation of [σ] of 0.5984) fit the data best (Figure 4). The Anderson-Darling Goodness of Fit test statistic (A^2) was 0.436 (P-value > 0.10). The equation of the fitted log-Gompertz model is of the form:

$$y = \Phi \left(\frac{x - 1.1443}{0.5984} \right)$$

Where x is the log (concentration) and y is the proportion of species affected, and Φ is the symbol representing normal distribution. Summary statistics for the long-term SSD are presented in Table 8-8. The 5th percentile on the short-term SSD is 1.5 mg/L. The lower fiducial limit (5%) on the 5th percentile is 1.2 mg/L, and the upper fiducial limit (95%) on the 5th percentile is 1.7 mg/L. The CWQGPAqL is defined as the 5th percentile on the SSD.

The long-term exposure CWQG for the protection of freshwater life is 1.5 mg/L.

Table 8- 8 Long-term CWQG for Boron Resulting from the SSD Method

	Concentration
SSD 5th percentile	1.5 mg/L
Lower 95% confidence limit	1.2 mg/L
Upper 95% confidence limit	1.7 mg/L




Figure 4 SSD for boron in freshwater derived by fitting the log-Normal model to the logarithm of acceptable long-term no- and low-effect endpoints for twenty-eight aquatic species versus Hazen plotting position (proportion of species affected)

Concentrations of boron in Canadian surface waters ranged from 0 to 2.58 mg/L total, 0.0001 to 0.951 mg/L extractable, and 0.0001 to 2.71 mg/L dissolved (Table 2-6). Total boron concentrations were not available for Nova Scotia, however, the maximum measured extractable boron concentration was 0.951 mg/L, which came from one river sample. This sample does not exceed the long-term WQGPAqL of 1.5 mg total boron / L. In the case of Nova Scotia, freshwater surface waters may be under the influence of marine water, which is higher in boron. Total boron concentrations exceeded the long-term WQGPAqL of 1.5 mg/L in Saskatchewan (maximum total boron measured was 2.58 mg/L) and in the Northwest Territories (maximum total boron measured was 2.3 mg/L). In the case of Saskatchewan and the Northwest Territories, these areas may be under the influence of natural boron mineral deposits. No measured surface water concentrations exceeded the short-term WQGPAqL of 29 mg/L.

In order to prevent adverse health effects to organisms caused by a deficiency of essential chemicals, recommended threshold levels for boron should not fall below the level required by the organism to remain healthy. Rowe *et al.* (1998) found that rainbow trout and zebrafish embryos experienced adverse effects at boron concentrations below 9.7×10^{-2} mg/L and 2.2×10^{-3} mg/L, respectively. Therefore, the proposed short-term benchmark concentration and long-term WQGPAqL meet the minimum boron requirements for both species.

9.0 RESEARCH NEEDS

The toxicity of many elements depends on the chemical speciation of the dissolved species. Boron was added as several water soluble forms for aquatic toxicity studies such as boric acid, sodium tetraborate, sodium borate, borax, sodium metaborate and sodium perborate. Although there were no statistically significant differences between the aquatic toxicity of boric acid and borax to aquatic species (Birge and Black, 1977), this may not hold true for other water soluble forms of boron. For example, metaborate releases hydrogen peroxide upon mixing with water, so the aquatic toxicity of this form of boron is known to be different (Guhl 1992) than the more environmentally relevant forms of boron, being undissociated boric acid and borate. Since the pKa of boric acid is 9.2, boric acid is the predominant form of boron in aquatic environments. Additional studies may be conducted to determine if the form of boron utilized during aquatic toxicity testing affects the toxicity of boron.

Several studies have shown that the low-level effects observed in reconstituted laboratory water are not predictive of the much higher effect levels observed under natural water exposure conditions (Butterwick *et al.*, 1989; Black *et al.*, 1993). These variations may be a result of differences in the natural chemical composition of waters from different geographical regions which may modify the toxicity of boron (Black *et al.*, 1993). Further studies investigating modifiers of boron aquatic toxicity are recommended.

10.0 OBJECTIVES OF OTHER AGENCIES

Moss and Nagpal (2003) report that the South African national boron criterion for coldwater-adapted species is 0.01 mg/L for acute effects (acute effect value or AEV) and 0.001 mg/L for chronic effects (chronic effect value or CEV) (Roux *et al.*, 1996). However, these values were not identified in the South African water quality guidelines (1996). South Africa established livestock watering and irrigation guidelines for boron of 0 to 5 mg/L and 0 to 0.5 mg/L, respectively (Republic of South Africa, 1996).

Australia and New Zealand adopted a freshwater high reliability trigger value for boron of 0.37 mg/L using the statistical distribution method at 95% protection (ANZECC, 2000). This was calculated using screened chronic freshwater data (around 30 points) from five taxonomic groups. Aquatic toxicity was expressed as NOEC equivalents or were adjusted to NOECs by dividing by factors depending on the endpoints (NOEC = MATC/2, LOEC/2.5 or E(L)C50/5) (ANZECC, 2000).

U.S. EPA (2006) does not provide a national ambient water quality criterion for freshwater aquatic life for boron, beyond referring back to the 1986 criterion which was based on irrigation of sensitive crops. Some individual states in the U.S. have set boron guidelines for freshwater aquatic life. Missouri has an effluent limit of 2 mg/L for subsurface waters (aquifer) for aquatic life protection (U.S. EPA, 1988). Missouri does not have a criterion for the protection of aquatic life; existing criteria only address irrigation and groundwater protection (Missouri CSR, 2005). New York State has set a chronic water quality standard for the protection of freshwater aquatic life of 10 mg/L (NYCRR, 1999). Region IV of the U.S. EPA has developed a chronic surface water screening benchmark of 0.75 mg/L (U.S. EPA Region IV, 2001). In addition, Region VI of the U.S. EPA recommended a freshwater surface water benchmark screening value of 0.0016 mg/L (TNRCC, 2001). Indiana Department of Environmental Management has developed both acute and chronic water screening benchmarks of 3.2 and 0.36 mg/L, respectively (USEPA GLI Scientific Criteria Document for the Development of a Canadian Water Quality Guideline for Boron

Clearinghouse, 2007). Ohio EPA has also developed both acute and chronic water screening benchmarks of 8.5 and 0.95 mg/L, respectively (USEPA GLI Clearinghouse, 2007).

CCME (1999) recommended a boron concentration of 5.0 mg/L for livestock drinking water. A boron concentration between 0.5 to 6 mg/L is recommended for agricultural irrigation water (CCME, 1999).

The Quebec Ministry of the Environment (Quebec Ministère du Développement durable, de l'Environnement et des Parcs) has boron freshwater guidelines for the protection of aquatic life that have been adopted from the Michigan Department of Environmental Quality. The Final Acute Value is 55 mg/L, the Acute Guideline (Final Acute Value divided by 2) is 28 mg/L, and the Chronic Guideline is 5 mg/L (MDDEP, 2008, pers. com.).

The maximum acceptable concentration of boron in Canadian drinking water was set to 5 mg/L (Health Canada, 1990). This interim maximum acceptable concentration was set as the availability of practicable treatment technology is inadequate to reduce boron concentrations in Canadian drinking water supplies to less than 5.0 mg/L in areas with high natural boron levels.

The British Columbia Water Protection Section of the Ministry of Water, Land and Air Protection has recommended that the boron concentration in freshwater not exceed 1.2 mg B/L for the protection of aquatic life (Moss and Nagpal, 2003). This guideline was derived based on a lowest effect level for growth inhibition on the green algae (*Selenastrum capricornutum*) of 12.3 mg/L (MELP, 1997). This lowest effect concentration was from a study which was chronic in duration and produced primary data. A safety factor of 0.1 was applied to derive the interim guideline (Moss and Nagpal, 2003).

11.0 REFERENCES

- Aldenberg, T., Jaworska, J.S., and Traas, T.P. 2002. Normal species sensitivity distributions and probabilistic ecological risk assessment. In: L. Posthuma, G.W. Suter II and T.P. Traas (Eds.), *Species Sensitivity Distributions in Ecotoxicology*. CRC Press LLC, Boca Raton, Florida. pp. 49-102.
- ANZECC. 2000. Australia and New Zealand Environment and Conservation Council and the Agriculture and Resource Management Council of Australia and New Zealand. *Australian and New Zealand Guidelines for Fresh and Marine Water Quality*. Volumes 1, 2 & 3.
- ATSDR. 1992. *Toxicological Profile for Boron and Compounds*. Agency for Toxic Substances and Disease Registry, U.S. Public Health Service. Available at: <http://www.atsdr.cdc.gov/toxprofiles/tp26.pdf>. [March 1, 2007].
- Avetisyan, L.T. 1983. Effectiveness of trace element utilisation in the fertilisation of piscicultural ponds of Armenia. *Biol. Zh. Arm.* 36: 22-29. *Cited In: Moss and Nagpal, 2003.*
- Bailer, J.A. and Oris, J.T. 1997. Estimating inhibition concentrations for different response scales using generalized linear models. *Environmental Toxicology and Chemistry* 16: 1554-1559.
- Bakke, J.P. 1991. Evaluation of the potential of boric acid to induce unscheduled DNA synthesis in the *in vitro* hepatocyte DNA repair assay using the male F-344 rat. (Unpublished study) Submitted by U.S. Borax Corp. MRID No. 42038903. *Cited In: U.S. EPA, 2004.*
- Benson, W.H., Birge, W.J., and Dorough, H.W. 1984. Absence of mutagenic activity of sodium borate (borax) and boric acid in the *Salmonella* pre-incubation test. *Environ. Toxicol. Chem.* 3: 209-214.
- Bergmann, W., Bruchlos, P. and Marks, G. 1995. The Toxic Limiting Value of Boron. *Tenside Sufactants Detergents.* 32(3):229-237.
- Bertine, K.K., and Goldberg, E.D. 1971. Fossil fuel combustion and the major sedimentary cycle. *Science* 173: 233-235.
- Bingham, E., Cohrssen, B., and Powell, C.H. 2001. *Patty's Toxicology 5th Edition*. Volume 3. John Wiley & Sons. New York, NY, p. 551.
- Birge, W.J. and Black, J.A. 1977. Sensitivity of vertebrate embryos to boron compounds. EPA-560/1-76-008. Office of Toxic Substances, USEPA, Washington, D.C.
- Birge, W.J. and Black, J.A. 1981. Toxicity of boron to embryonic and larval stages of Largemouth Bass (*Micropterus salmoides*) and Rainbow Trout (*Salmo gairdneri*). Completion report prepared for Proctor & Gamble, Unpublished. *Cited In: Butterwick et al., 1989, Cited In: Moss and Nagpal, 2003.*
- Birge, W.J., J.A. Black, A.G. Westerman and B.A. Ramey. 1983. Fish and Amphibian Embryos - A Model System for Evaluating Teratogenicity. *Fundamental and Applied Toxicology.* 3:237-242.
- Black, J.A., Barnum, J.B. and Birge, W.J. 1993. An integrated assessment of the effects of boron to the rainbow trout. *Chemosphere* 26: 1383-1413.
- Borgmann, U., Couillard, Y., Doyle, P., and Dixon, G.D. 2005. Toxicity of sixty-three metals and metalloids to *Hyalella azteca* at two levels of water hardness. *Environmental Toxicology and Chemistry* 24(3): 641-652.
- Bringmann, G. 1978. Determining the harmful effect of water pollutants on protozoa. 1. Bacteriovorous flagellates. *Z. Wasser Abwasser Forsch.* 11: 210-215.

- Budavari, S., O'Neil, M.J., Smith, A., Heckelman, P.E. 1989. The Merck Index: an encyclopedia of chemicals, drugs, and biologicals. Merck & Co., Inc, Rahway, N.J, pp. 1349.
- Butterwick, L., De Oude, N., and Raymond, K. 1989. Safety assessment of boron in aquatic and terrestrial Environments. *Ecotoxicology and Environmental Safety*. 17: 339-371.
- CCME (Canadian Council of Ministers of the Environment). 1991. Appendix IX— A protocol for the derivation of water quality guidelines for the protection of aquatic life (April 1991). In: Canadian water quality guidelines, Canadian Council of Resource and Environment Ministers, 1987. Prepared by the Task Force on Water Quality Guidelines. [Updated and reprinted with minor revisions and editorial changes in Canadian environmental quality guidelines, Chapter 4, Canadian Council of Ministers of the Environment, 1999, Winnipeg.]
- CCME. 1993. Protocols for Deriving Water Quality Guidelines for the Protection of Agricultural Water Uses. Appendix XV. Canadian Council of Ministers of the Environment.
- CCME. 1999. Canadian water quality guidelines for the protection of agricultural water uses: Summary table. In: Canadian environmental quality guidelines, 1999, Canadian Council of Ministers of the Environment, Winnipeg.
- CCME. 2007. A Protocol for the Derivation of Water Quality Guidelines for the Protection of Aquatic Life. Canadian Council of Ministers of the Environment. Draft for Public Review. April 27, 2007.
- CESARS. 1999. Boron. Chemical Evaluation Search and Retrieval System, Canadian Centre for Occupational Health and Safety. Available at: <http://ccinfoweb.ccohs.ca/chempendium/search.html>. [February 13, 2007].
- CHEMINFO. 1996. Boric acid. CHEMINFO, Canadian Centre for Occupational Health and Safety. Available at: <http://ccinfoweb.ccohs.ca/chempendium/search.html>. [March 2, 2007].
- Clayton, G. D., and Clayton, F.E. (Eds.). 1982. Patty's Industrial Hygiene and Toxicology 3rd Edition. Volume 2a-c. John Wiley and Sons, New York, NY.
- Davis, J.C. and Mason, B.J. 1973. Toxicity of Boron to Pacific Salmon, and Boron Uptake Experiments with Pacific Salmon and Oysters. Appendix III. In *Zinc and Boron Pollution in Coastal Waters of British Columbia by Effluents from the Pulp and Paper Industry*. 1973. Environment Canada, Pacific Region, Vancouver, B.C.
- Davis, S.M., Drake, K.D., and Maier, K.J. 2002. Toxicity of boron to duckweed, *Spirodella polyrrhiza*. *Chemosphere* 48: 615-620.
- Demerec, M., Bentani, G., and Flint, J. 1951. A survey of chemicals for mutagenic action on *E. coli*. *Am Nat* 84(821): 119-136. *Cited In: U.S. EPA, 2004.*
- Durocher, N.L. 1969. Preliminary air pollution survey of boron and its compounds. A literature review prepared under Contract No. PH 22-68-25, Public Health Service, National Air Pollution Control Administration, U.S. Department of Health, Education and Welfare, Raleigh, NC. *Cited In: Health Canada, 1990.*
- Dyer, S.D. 2001. Determination of the aquatic PNEC_{0.05} for boron. *Chemosphere*. 44:369-376.
- Eisler, R. 1990. Boron hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Fish Wildlife Service., Biological Report 85(1.20). 32pp.
- Fay, R.W. 1959. Toxic effects of boron trioxide against immature stages of *Aedes aegypti*, *Anopheles quadrimaculatus* and *Cuex quinquefasciatus*. *J. Econ. Entomol.* 52: 1027- 1028.
- Fernandez, E., Sanchez, E., Bonilla, I., Mateo, P., and Ortega, P. 1984. Effect of boron on the growth and cell composition of *Chlorella pyrenoidosa*. *Phyton*. 44: 125-131.

- Fort, D.J., Propst, T.L., Stover, E.L., Murray, F.J., and Strong, P.L. 1999. Adverse effects for low dietary and environmental boron exposure on reproduction, development and maturation in *Xenopus laevis*. The Journal of Trace Elements in Experimental Medicine 12: 175-185.
- Frick, H. 1985. Boron tolerance and accumulation in the duckweed, *Lemna minor*. J. Plant. Nut. 8(12): 1123-1129.
- Gersich, F.M. 1984. Evaluation of a static renewal chronic toxicity test method for *Daphnia magna* Straus using boric acid. Environ. Toxicol. Chem. 3: 89-94.
- Glandon, R.P., and McNabb, C.D. 1978. The uptake of boron by *Lemna minor*. Aquatic Botany 4: 53-64.
- Guhl, W., 1992a. Okologischespektrum von bor. SOFW J. 118 (18/92), 1159-1168. Cited In: Dyer, S.D. 2001.
- Guhl, W., 1992b. Laboratory river models and their relevance to the real environment. Presentation at Joint Meeting of SETAC-Europe, Potsdam, Germany, 22-24 June 1992. Cited In: Dyer, S.D. 2001.
- Hamilton, S.J. 1995. Hazard assessment of inorganics to three endangered fish in the Green River, Utah. Ecotoxicology and Environmental Safety 30: 134-142
- Hamilton, S.J. and Buhl, K.J. 1990. Acute toxicity of boron, molybdenum, and selenium to fry of chinook salmon and coho salmon. Arch. Environ. Contam. Toxicol. 19: 366-373.
- Hawley, G.G. 1981. The Condensed Chemical Dictionary 10th Edition. Van Nostrand Reinhold Co., New York, NY. Cited In: CESARS, 1999.
- Haworth, S., Lawlor, T., Mortelmans, K., Speck, W., and Zeiger, E. 1983. *Salmonella* mutagenicity test results for 250 chemicals. Environ Mutagen, 1(Suppl): 3-142. Cited In: WHO, 1998.
- Health Canada. 1990. Guidelines for Canadian Drinking Water Quality: Boron. Available at: http://www.hc-sc.gc.ca/ewh-semt/pubs/water-eau/doc_sup-appui/boron-bore/index_e.html. [February 27, 2007].
- Hickey, C.W. 1989. Sensitivity of four New Zealand cladoceran species and *Daphnia magna* to aquatic toxicants. N.Z. J. Marine Freshwater Res. 23,131-137.
- Hooftman et al. 2000. TNO Nutrition and Food Research Institute, The Netherlands. TNO Report V99-168 Unpublished Report to Borax Europe, Ltd. Cited In: Neobor® (*Borax Pentahydrate*) MSDS, U.S. Borax Inc.,
- Howe, P.D. 1998. A review of boron effects in the environment. Biological Trace Element Research 66: 153-166.
- HSDB. 2005. Hazardous Substances Databank. Boron. Available at: <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>. [March 22, 2007].
- Hydromantis Inc., Minnow Environmental Inc., and University of Waterloo. 2005. Review of the State of Knowledge of Municipal Effluent Science and Research – Review of Effluent Substances. Report Prepared for: Development Committee for the MWWWE Canada-Wide Strategy Canadian Council of Ministers of the Environment. PN 1356, June 2005.
- Iyer, V.N., and Szybalski, W. 1958. Two simple methods for the detection of chemical mutagens. Appl Microbiol 6: 23-29.
- Kapu, M.M. and Schaeffer, D.J. 1991. Planarians in Toxicology. Responses of asexual *Dugesia dorotocephala* to selected metals. Bull. Environ. Contam. Toxicol. 47: 302- 307.
- Kerr, D.R. and Meador, J.P. 1996. Modeling dose response using generalized linear models. Environmental Toxicology and Chemistry 15: 395-401.

- Kirk-Othmer. 1992. Kirk-Othmer Encyclopedia of Chemical Technology 4th Edition. Volume 4. Jon Wiley and Sons, New York, NY, p. 375. *Cited In: HSDB, 2005 and CHEMINFO, 1996.*
- Landolph, J.R. 1985. Cytotoxicity and negligible genotoxicity of borax ores to cultured mammalian cells. *Am. J. Ind. Med.* 7: 31-43. *Cited In: Eisler, 1990 and WHO, 1998.*
- Laposata, M.M., and Dunson, W.A. 1998. Effects of boron and nitrate on hatching success of amphibian eggs. *Arch. Environ. Contam. Toxicol* 35: 615-619.
- Laws, E.A. 1981. Aquatic Pollution. John Wiley and Sons, NY. *Cited In: Moss and Nagpal, 2003.*
- Lewis, M.A. and Valentine, L.C. 1981. Acute and chronic toxicities of boric acid to *Daphnia magna* Straus. *Bull. Environ. Contam. Toxicol.* 27: 309-315.
- Lewis, R.J. 1999. Sax's Dangerous Properties of Industrial Materials 10th Edition. Volumes 1-3. John Wiley & Sons Inc., New York, NY. *Cited In: HSDB, 2005.*
- Lide, D.R. (Ed.). 2000. CRC Handbook of Chemistry and Physics 81st Edition. CRC Press LLC, Boca Raton, Florida. *Cited In: HSDB, 2005.*
- Lochner, C. 2008. Water Quality Monitoring and Surveillance, Water Science and Technology, Environment Canada. Boron in Canadian Fresh Waters. Personal Communication.
- Maeso, E.S., Valiente, E.F., Bonilla, I. and Mateo, P. 1985. Accumulation of proteins in giant cells, induced by high boron concentrations in *Chlorella pyrenoidosa*. *J. Plant Physiol.* 121: 301-311.
- Maier, K.J. and Knight, A.W. 1991. The toxicity of waterborne boron to *Daphnia magna* and *Chironomus decorus* and the effects of water hardness and sulfate on boron Toxicity. *Arch. Environ. Contam. Toxicol.* 20: 282-287.
- Martinez, F., Mateo, P., Bonilla, I., Fernandez-Valiente, E., and Garate, A. 1986. Growth of *Anacystis nidulans* in relation to boron supply. *Irs. J. Bot.* 35: 17-21.
- Mateo, P., Martinez, F., Bonilla, I., Valiente, E.F., and Maeso, E.S. 1987. Effects of high boron concentrations on nitrate utilization and photosynthesis in blue-green algae *Anabaena* PCC 7119 and *Anacystis nidulans*. *J. Plant Physiol.* 128: 161-168.
- McGregor, D.B., Brown, A., Cattnach, P., Edwards, I., McBride, D., Riach, C., and Caspari, W.J. 1988. Responses of the L5178Y tk-/tk- mouse lymphoma cell forward mutation assay: III. 72 coded chemicals. *Environ Mol Mutagen* 12: 85-154. *Cited In: WHO, 1998.*
- McKee, J.E. and Wolf, H.W. 1963. Water Quality Criteria. The Resource Agency of California. 2nd Ed. State Water Quality Control Board. Public. No. 3A.
- MDDEP (Quebec Ministère du Développement durable, de l'Environnement et des Parcs). 2008. Unpublished boron water quality monitoring data in 23 rivers in Quebec. Ministère du Développement durable, de l'Environnement et des Parcs. S. Cloutier, personal communication.
- MELP (British Columbia Ministry of Environment, Lands and Parks). 1996. Unpublished data on boron toxicity on *Oncorhynchus kisutch*, *Hyalella Azteca*, *Eohaustorius washingtonianus* and Microtox. Ministry of Environment, Lands and Parks. *Cited In: Moss and Nagpal, 2003.*
- MELP. 1997. Unpublished data on boron toxicity on *Selenastrum capricornutum* and *Strongylocentrus droebachiensis*. Ministry of Environment, Lands and Parks. *Cited In: Moss and Nagpal, 2003.*
- Missouri CSR (Code of State Regulations). 2005. 10 CSR 20-7—Department of Natural Resources Division 20—Clean Water Commission. Table A—Criteria for Designated Uses(11/30/05). <http://www.sos.mo.gov/adrules/csr/current/10csr/10c20-7b.pdf>.

- MOE. 1992. Ontario's Water Quality Objective Development Process. Ontario Ministry of the Environment.
- Moore, D.R.J., and Caux, P.Y. 1997. Estimating low toxic effects. *Environmental Toxicology and Chemistry* 16(4): 794-801.
- Moss, S.A., and Nagpal, N.K. 2003. Ambient Water Quality Guidelines for Boron- Full Report. British Columbia. Water, Air and Climate Change Branch.
- NAS. 1973. Water Quality Criteria 1972. National Academy of Sciences, National Academy for Engineering. EPA Ecol. Res. Serv. US. Environmental Protection Agency Rep. R3-73-033.
- Newman, M.C., Ownby, D.R., Mézin, L.C.A., Powell, D.C., Christensen, T.R.L., Lerberg, S.B., Anderson, B-A., and Padma, T.V. 2002. Species sensitivity distributions in ecological risk assessment: Distributional assumptions, alternate bootstrap techniques, and estimation of adequate number of species. In: Posthuma, L., G.W. Suter II and T.P. Traas (Eds.). *Species Sensitivity Distributions in Ecotoxicology*. Lewis Publishers, New York, NY. pp119-132.
- NIOSH. 2003. NIOSH Pocket Guide to Chemical Hazards: Borates, tetra, sodium salts (Decahydrate). The National Institute for Occupational Safety and Health.
- Nobel, W., Mayer, T., and Kohler, A. 1983. Submerged water plants as testing organisms for pollutants. *Zeitschrift für Wasser und Abwasser Forschung*. 16(3): 87-90.
- NTP. 1987. Toxicology and carcinogenesis studies of boric acid (CAS No. 10043-35-3) in B6C3F₁ mice (feed studies). Research Triangle Park, North Carolina, US Department of Health and Human Services, National Toxicology Program (NTP TR No. 324).
- NYCRR (New York Codes, Rules and Regulations). 1999. Surface Water and Groundwater Quality Standards and Groundwater Effluent Limitations. 6 NYCRR Part 703. §703.5 Table 1. Water Quality Standards for Surface Waters and Groundwater, Water quality standards for taste, color- and odor-producing, toxic and other deleterious substances. <http://www.dec.state.ny.us/website/regs/part703.html#table1>
- Oduola, O.A. 1997. Individual and combined genotoxic response of boric acid and aflatoxin B₁ in *Escherichia coli* PQ37. *East Afr Med. J* 74: 499-502. *Cited In: U.S. EPA, 2004.*
- O'Loughlin, K.G. 1991. Bone marrow erythrocyte micronucleus assay of boric acid in Swiss-Webster mice. (Unpublished study) Submitted by U.S. Borax Corp. MRID No. 42038904. *Cited In: U.S. EPA, 2004.*
- O'Neil, M.J. (Ed.). 2001. The Merck Index- An Encyclopedia of Chemicals, Drugs and Biologicals 13th Edition. Merck and Co., Inc., Whitehouse Station, N.J. *Cited In: HSDB, 2005.*
- Park, H. and Schlesinger, W.H. 2002. Global biogeochemical cycle of boron. *Global Biogeochemical Cycles*. 16(4):1-11.
- Proctor and Gamble. 1979. Unpublished data – Fathead minnow toxicity testing using boric acid. *Cited In: Moss and Nagpal, 2003.*
- Rai, D., *et al.* 1986. Chemical attenuation rates, coefficients, and constants in leachate migration. Volume 1: A critical review. Report to Electric Power Research Institute, Palo Alto, CA, by Battelle Pacific Northwest Laboratories, Richland, WA. Research Project 2198-1. *Cited In: WHO, 2003.*
- Republic of South Africa. 1996. South African Water Quality Guidelines 2nd Edition. Volume 1-8. Republic of South Africa, Department of Water Affairs and Forestry. Available at: http://www.dwaf.gov.za/IWQS/wq_guide/v.za/IWQS/wq_guide/agriaqua.pdf. [March 22, 2007].
- Rosenkranz, H.S. 1973. Sodium hypochlorite and sodium perborate: Preferential inhibitors of DNA polymerase-deficient bacteria. *Mutat Res*, 21: 171-174. *Cited In: WHO, 1998.*
- Rossmann, R., and Barres, J. 1988. Trace element concentrations in near-surface waters of the great lakes and methods of collection, storage and analysis. *J. Great Lakes Res.* 14(2): 188-204.

- Rowe, R.I., Bouzan, C., Nabili, S., and Eckhart, C.D. 1998. The response of trout and zebrafish embryos to low and high boron concentrations is u-shaped. *Biol. Trace Element Res* 66: 261-270.
- Roux, D.J., S.H.J. Jooste and H.M. MacKay. 1996. Substance-specific water quality criteria for the protection of South African freshwater ecosystems: methods for derivation and initial results for some inorganic toxic substances. *South African J. Science*. Vol 92. *Cited In: Moss and Nagpal, 2003.*
- Rudd, C.J. 1991. Mouse lymphoma cell mutagenesis assay (tK+/-/tK-/-) of boric acid. (Unpublished study) Submitted by U.S. Borax Corp. MRID No. 4203902. *Cited In: U.S. EPA, 2004.*
- Saiki, M.K., Jennings, M.R., and Brumbaugh, W.G. 1993. Boron, molybdenum, and selenium in aquatic food chains from the lower San Joaquin River and its tributaries, California. *Arch. Environ. Contam. Toxicol.* 24: 307-319.
- Sanchez, E.M. *et al.* 1982. Comportamiento de *Chlorella pyrenoidosa* frente a distintas concentraciones de boro en el medio. *Anal Edaf Agrobiol.* 41: 2357-2362. *Cited In: Maier and Knight, 1991, Cited In: Moss and Nagpal, 2003.*
- Sax, N.I. 1984. *Dangerous Properties of Industrial Materials* 6th Edition. Van Nostrand Reinhold Co., New York, NY. *Cited In: CESARS, 1999.*
- Sekerka, I., and Lechner, J.F. 1990. Automated method for the determination of boron in water by flow-injection analysis with in-line preconcentration and spectrophotometric detection. *Analytica Chimica Acta* 234: 199-206.
- Shiu, W.Y., *et al.* 1990. *Rev Environ Contam Toxicol* 116: 15-187. *Cited In: HSDB, 2005.*
- Sprague, R.W. 1972. The ecological significance of boron. United States Borax and Chemical Corp., Los Angeles. 58pp.
- Stanley, R.A. 1974. Toxicity of heavy metals and salts to Eurasian watermill (*Myriophyllum spicatum*). *Arch. Environ. Contam. Toxicol.* 2: 331-341.
- Stewart, K.R. 1991. *Salmonella*/microsome plate incorporation assay of boric acid. (Unpublished study) Submitted by U.S. Borax Corp. MRID No. 4203901. *Cited In: U.S. EPA, 2004.*
- Stokinger, H.E. 1981. Boron. In: Clayton GD, Clayton FE, eds. *Patty's industrial hygiene and toxicology*. Vol. 2B. Toxicology. 3rd ed. New York, NY: John Wiley and Sons, 2978-3005. *Cited In: ATSDR, 1992.*
- Suloway, J.J., Roy, W.R., Skelly, T.M., Dickerson, D.R., Schuller, R.M., and Griffin, R.A. 1983. Chemical and toxicological properties of coal fly ash. Champaign, Illinois, Illinois State Geological Survey (Publication NTIS PB84-116110). *Cited In: WHO, 1998.*
- Szybalski, W. 1958. Special microbiological system. II. Observations of chemical mutagenesis in microorganisms. *Ann NY Acad. Sci* 76: 475-489. *Cited In: U.S. EPA, 2004.*
- Takeuchi, T. 1958. Effects of boric acid on the development of the eggs of the toad, *Bufo vulgaris formosus*. *Sci. Rep. Tohoku Univ. Ser. 4 Biol.* 24: 33-43.
- Thompson, S.E., Burton, C.A., Quinn, D.J., *et al.* 1972. Concentration factors of chemical elements in edible aquatic organisms. Report to U.S. Atomic Energy Commission by Lawrence Livermore Laboratory, University of California, Livermore, CA. UCRL-50564. *Cited In: ATSDR, 1992.*
- Thompson, J.A.J., Davis, J.C. and Drew, R.E. 1976. Toxicity, uptake and survey studies of boron in the marine environment. *Water Res.* 10: 869-875.
- TNRCC. 2001. Guidance for Conducting Ecological Risk Assessment at Remediation Sites in Texas. Toxicology and Risk Assessment Section, Texas Natural Resource Conservation Commission, Austin, TX. RG-263.

- Tsui, P.T., and McCart, P.J. 1981. Chlorinated hydrocarbon residues and heavy metals in several fish species from the Cold Lake area in Alberta, Canada. *Intern J Environ Anal Chem* 10: 277-285. *Cited In: ATSDR, 1992.*
- Turnbull, H. *et al.* 1954. Toxicity of various refinery materials to freshwater fish. *Ind. Eng. Chem*, 46: 324-333.
- U.S. Bureau of Mines. 1989. Mineral commodity summaries. Washington, DC. *Cited In: ATSDR, 1992.*
- U.S. EPA. 1975. Preliminary investigation of effects on the environment of boron, indium, nickel, selenium, tin, vanadium and their compounds. Vol. 1. Boron. U.S. Environmental Protection Agency Rep. 56/2-75-005A. 111pp. *Cited In: Eisler, 1990.*
- U.S. EPA. 1987. Health effects assessment for boron and compounds. Environmental Criteria and Assessment Office, U.S. Environmental Protection Agency. *Cited In: Health Canada, 1990.*
- U.S. EPA. 1988. Office of Water. Water Quality Standards Criteria Summaries: A Compilation of State/Federal Criteria. EPA/440/5-88/011.
- U.S. EPA Region IV. 2001. Supplemental Guidance to RAGS: Region 4 Bulletins, Ecological Risk Assessment. Available at: <http://www.epa.gov/region4/waste/ots/ecolbul.htm>. [March 22, 2007].
- U.S. EPA. 2004. Carcinogenicity Assessment: Boron and Compounds (CASRN 7440-42-8). Integrated Risk Information System (IRIS), U.S. Environmental Protection Agency. Available at: <http://www.epa.gov/iris/subst/0410.htm>. [March 1, 2007].
- U.S. EPA. 2006. National Recommended Water Quality Criteria. Office of Water. Office of Science and Technology (4303T). U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 2007. Great Lakes Initiative (GLI) Toxicity Data Clearinghouse. Water Quality Branch, US EPA Regions 2, 3, and 5, and the Great Lakes States and Tribes. Available at: http://cfpub.epa.gov/gliclear/criteria_query.cfm [May 10, 2007].
- Wallen, I.E. *et al.* 1957. Toxicity to *Gambusia affinis* of certain pure chemicals in turbid water. *Sewage Ind. Wastes* 29: 695-711.
- Wang, W. 1986. Toxicity Tests of Aquatic Pollutants by Using Common Duckweed. *Environ. Poll. (Series B)* 11:1-14.
- Weast, R.C. (Ed). 1985. CRC handbook of chemistry and physics. Boca Raton, FL: CRC Press, Inc., B-79, B-10. *Cited In: ATSDR, 1992 and WHO, 2003.*
- WHO. 1998. Environmental Health Criteria Document No. 204: Boron (7740-42-8). World Health Organization.
- WHO. 2003. Background document for development of WHO Guidelines for Drinking-water Quality: Boron in Drinking-water. World Health Organization.
- Windholz, M. (Ed). 1983. The Merck index: An encyclopedia of chemicals, drugs, and biologicals. 10th ed. Rahway, NJ: Merck and Company, Inc. 186-187. *Cited In: ATSDR, 1992.*
- Wren, C.D., MacCrimmon, H.R., and Loescher, B.R. 1983. Examination of bioaccumulation and biomagnification of metals in a Precambrian shield lake. *Water Air Soil Pollut* 19: 277-291.
- Wurtz, A. 1945. The action of boric acid on certain fish: trout, roach, rudd. *Ann. Stn. Cent. Hydrobiol. Appl.* 1: 179.

APPENDIX A

AQUATIC TOXICITY TABLE FOR BORON

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
VERTEBRATES – Acute													
Largemouth bass	<i>Micropterus salmoides</i>	embryo-hatching stages	Teratogenesis at Hatching (3 d, NOEC)	7.5	20	8.4		204	0.109	A,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Largemouth bass	<i>Micropterus salmoides</i>	embryo-hatching stages	Teratogenesis at Hatching (3 d, LOEC)	7.5	20	8.4		204	1.39	A,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Largemouth bass	<i>Micropterus salmoides</i>	embryo-hatching stages	Mortality at Hatching (3 d, NOEC)	7.5	20	8.4		204	1.39	A,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Largemouth bass	<i>Micropterus salmoides</i>	embryo-hatching stages	Mortality at Hatching (3 d, LOEC)	7.5	20	8.4		204	12.17	A,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Bluegill sunfish	<i>Lepomis macrochirus</i>	average size 7 cm, 5 g	Mortality (24 hr, TLm)	6.9-7.5	20		33-81	84-163	4.6	A,S	S	Turnbull <i>et al.</i> , 1954	Sodium tetraborate
Bluegill sunfish	<i>Lepomis macrochirus</i>	average size 7 cm, 5 g	Mortality (24 hr, TLm)		20				2,389	A,S	S	Turnbull <i>et al.</i> , 1954	Boron trifluoride
Fathead Minnow	<i>Pimephales promelas</i>	Larval	96-h LC50	7.5-7.9	18-22	5-8.4	85-90	120	64.3	A,S,M	P	MOE 2007 (unpublished data)	Boric acid
Fathead Minnow	<i>Pimephales promelas</i>	larval	48-h LC50	7.5-7.9	18-22	5-8.4	85-90	120	348	A,S,M	P	MOE 2007 (unpublished data)	Boric acid
Zebrafish	<i>Danio rerio</i>	(zygotes) fertilization to free feeding	Embryonic Mortality (96 hr, LOEC, 88% mortality)						99.5 (9.2 mmol/L) ^a	A,S,M	S	Rowe <i>et al.</i> , 1998	Boric acid
Bonytail	<i>Gila elegans</i>	1.1 g juvenile	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	>100	A,S,U	S	Hamilton, 1995	Boric acid
Bonytail	<i>Gila elegans</i>	swim-up fry	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	280	A,S,U	S	Hamilton, 1995	Boric acid
Bonytail	<i>Gila elegans</i>	2.6 g juvenile	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	552	A,S,U	S	Hamilton, 1995	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
Colorado squawfish	<i>Ptychocheilus lucius</i>	0.4-1.1g juvenile	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	>100	A,S,U	S	Hamilton, 1995	Boric acid
Colorado squawfish	<i>Ptychocheilus lucius</i>	swim-up fry	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	279	A,S,U	S	Hamilton, 1995	Boric acid
Colorado squawfish	<i>Ptychocheilus lucius</i>	1.7 g juvenile	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	527	A,S,U	S	Hamilton, 1995	Boric acid
Razorback sucker	<i>Xyrauchen texanus</i>	2.0 g juvenile	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	>100	A,S,U	S	Hamilton, 1995	Boric acid
Razorback sucker	<i>Xyrauchen texanus</i>	swim-up fry	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	233	A,S,U	S	Hamilton, 1995	Boric acid
Razorback sucker	<i>Xyrauchen texanus</i>	0.9 g juvenile	Mortality (96 hr, LC50)	7.0-8.5	24-26	>40% saturation	105-109	191-201	279	A,S,U	S	Hamilton, 1995	Boric acid
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (96 hr, NOEC)	8.6-9.1	22-26				<204	A,S	S	Wallen <i>et al.</i> , 1957	Sodium tetraborate
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (144 hr, TLm)	8.6-9.1	22-26				215	A,S	S	Wallen <i>et al.</i> , 1957	Sodium tetraborate
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (96 hr, NOEC)	5.4-7.3	20-23				<314	A,S	S	Wallen <i>et al.</i> , 1957	Boric acid
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (96 hr, TLm)	8.6-9.1	22-26				408	A,S	S	Wallen <i>et al.</i> , 1957	Sodium tetraborate
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (48 hr, TLm)	8.6-9.1	22-26				929	A,S	S	Wallen <i>et al.</i> , 1957	Sodium tetraborate
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (96 hr, TLm)	5.4-7.3	20-23				979	A	S	Wallen <i>et al.</i> , 1957	Boric acid
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (24 hr, TLm)	8.6-9.1	22-26				1,360	A,S	S	Wallen <i>et al.</i> , 1957	Sodium tetraborate
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (48 hr, TLm)	5.4-7.3	20-23				1,834	A,S	S	Wallen <i>et al.</i> , 1957	Boric acid
Mosquito fish	<i>Gambusia affinis</i>	adult females	Mortality (24 hr, TLm)	5.4-7.3	20-23				3,145	A,S	S	Wallen <i>et al.</i> , 1957	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	early life stages	Mortality (96hr, LC50)	7.5-7.9	14-16	5-8.4	85-90	120	275	A,S,M	P	MOE 2007 (unpublished data)	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	early life stages	Mortality (96hr, LC50)		14-16			250	336*	A	P*	MELP, 1996	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	adults	Mortality (48 hr, LC50)						339	A	U	Sprague, 1972	
Rainbow trout	<i>Oncorhynchus</i>	adults	(0.5 hr, NOEC)						350	A	U	Sprague,	

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
	<i>mykiss</i>										1972		
Rainbow trout	<i>Oncorhynchus mykiss</i>	early life stages	Mortality (96hr, LC50)		14-16			100	379.6	A	P*	MELP, 1996	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	early life stages	Mortality (96hr, LC50)		14-16			25	436.2	A	P*	MELP, 1996	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	early life stages	(7 d, EC50-embryo test)						969	A,R,	P*	MELP, 1996	
Rainbow trout	<i>Oncorhynchus mykiss</i>	adults	Obvious distress (0.5 hr)						3,500	A	U	Sprague, 1972	
Coho salmon	<i>Oncorhynchus kisutch</i>	salmonids	Mortality (96 hr, LC50)		14-16			100	304.1	A	P*	MELP, 1996	Boric acid
Coho salmon	<i>Oncorhynchus kisutch</i>	salmonids	Mortality (96 hr, LC50)		14-16			25	357.4	A	P*	MELP, 1996	Boric acid
Coho salmon	<i>Oncorhynchus kisutch</i>	swim-up fry, 0.5 g mean weight	Mortality (96 hr, LC50)	7.47-8.17	12			210-212	447	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Coho salmon	<i>Oncorhynchus kisutch</i>	salmonids	Mortality (96 hr, LC50)		14-16			250	477.1	A	P*	MELP, 1996	Boric acid
Coho salmon	<i>Oncorhynchus kisutch</i>	swim-up fry, 0.5 g mean weight	Mortality (24 hr, LC50)	7.47-8.17	12			210-212	>1,000	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Minnow			Mortality (minimum lethal dose)		19				340-374	A	?	LeClerc and Devlaminck (1950, 1955) cited in McKee and Wolf, 1963	Sodium tetraborate
Minnow			Mortality (minimum lethal dose)		17			hard	793-850	A	?	LeClerc and Devlaminck (1950, 1955) cited in McKee and Wolf, 1963	Sodium tetraborate
Minnow		between 5-8 cm	Mortality (6 hr, minimum lethal dose or LOEC)		20	6.42	12.5	Distilled	3,145-3,319	A	U	LeClerc, 1960	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
Minnow		between 5-8 cm	Mortality (6 hr, minimum lethal dose or LOEC)		20	6.42	150	hard	3,319-3,407	A	U	LeClerc, 1960	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	0.31 g	Mortality (96hr, LC50)	7.51-7.63	12			41.4-42.0	566	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	swim-up fry, mean weight 1.1g	Mortality (96hr, LC50)	7.47-8.17	12			210-212	725	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	swim-up fry, mean weight 1.1g	Mortality (24 hr, LC50)	7.47-8.17	12			210-212	>1,000	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	eyed egg	Mortality (24 hr, LC50)	7.51-7.63	12			41.4-42.0	>1,000	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	eyed egg	Mortality (96hr, LC50)	7.51-7.63	12			41.4-42.0	>1,000	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	alevin	Mortality (24 hr, LC50)	7.51-7.63	12			41.4-42.0	>1,000	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	alevin	Mortality (96hr, LC50)	7.51-7.63	12			41.4-42.0	>1,000	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	0.31 g	Mortality (24 hr, LC50)	7.51-7.63	12			41.4-42.0	>1,000	A,S,U	S	Hamilton and Buhl, 1990	Boric acid
Frog	<i>Xenopus laevis</i>	embryo-larval	Abnormal Development (4 d, NOEC)	8	22-24				35.9	A,R,M	S	Fort <i>et al.</i> , 1999	Boric acid
Frog	<i>Xenopus laevis</i>	embryo-larval	Abnormal Development (4 d, LOEC)	8	22-24				53.85	A,R,M	S	Fort <i>et al.</i> , 1999	Boric acid
Toad	<i>Bufo vulgaris</i>	embryo, from 2 cell stage to tailbud stage	Malformation (24 hr, edema, microcephalia, short tail and suppressed forebrain development)						874	A	?	U.S. EPA, 1975 In Eisler, 1990	Boric acid
Toad	<i>Bufo vulgaris formosus</i>	embryo	Teratogenic Defects and Reduced Survival						1,747 (1% solution)	A,S	U	Takeuchi, 1958	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
			(24 hr)										
<u>INVERTEBRATES – Acute</u>													
Flatworm	<i>Dugesia dorocephala</i>	18-20 mm	Behaviour (1 hr, restlessness, hyperkinesia, spiralling and head/nose twist within 5 min. exposure)	6.0-8.0				370	1	A,S,U	U	Kapu and Schaeffer, 1991	Boron
Flatworm	<i>Dugesia dorocephala</i>	18-20 mm	Behaviour (1 hr, restlessness, hyperkinesia, spiralling and head/nose twist within 5 min. exposure)	6.0-8.0				370	10	A,S,U	U	Kapu and Schaeffer, 1991	Boron
Protozoan	<i>Entosiphon sulcatum</i>		Cell Replication (72 hr, TT, 5% reduction in cell replication)	6.9	25				1	A,S	? (English abstract, remainder in German)	Bringmann, 1978 (English abstract, remainder in German)	Sodium tetraborate
Amphipod	<i>Hyalella azteca</i>	1-11 d	Mortality (48h, LC50)	7.5-7.9	22	5-8.4	85-90	120	140	A,S,M	P	MOE, 2007 (unpublished data)	Boric acid
Amphipod	<i>Hyalella azteca</i>		Mortality (96 hr, LC50)		22-24			25	28.9	A	P*	MELP, 1996	Boric acid
Amphipod	<i>Hyalella azteca</i>		Mortality (96 hr, LC50)		22-24			100	291.3	A	P*	MELP, 1996	Boric acid
Amphipod	<i>Hyalella azteca</i>		Mortality (96 hr, LC50)		22-24			250	333.6	A	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i>	neonates	Mortality (48 hr, LC50)		19-21			25	21.3	A	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i>	< 24 hr old	(NOEC)					100	25.6	A,S	P*	MELP, 1996	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
Water Flea	<i>Daphnia magna</i>	< 24 hr old	Mortality (LOEC)					100	50	A,S	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i>	< 24 hr old	(NOEC)					250	50	A,S	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i>	neonates	Mortality (48 hr, LC50)		19-21			100	52.4	A	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i>	< 24 hr old	Mortality (LOEC)					250	100	A,S	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i> straus	< 24 hr old	Mortality (48 hr, LC50)	6.7-8.1	20.1-20.7	5.39 (> 60% saturation)	53-63	141-155	133 (115-153)	A,S,U	S	Gersich, 1984	Boric acid
Water Flea	<i>Daphnia magna</i>	neonates	Mortality (48 hr, LC50)		19-21			250	139.2	A	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i>	neonate	Mortality (48 hr, LC50)	9.0-9.2	19.9-20.1	8.4-8.8		10.6-170	141	A,R,M	P	Maier and Knight, 1991	Sodium tetraborate
Water Flea	<i>Daphnia magna</i>	<24 h old	Mortality (48-h LC50)	7.5-7.9	18-22	5-8.4	85-90	120	165	A,S,M	P	MOE 2007 (unpublished data)	Boric acid
Water Flea	<i>Daphnia magna</i> straus	<24 hr old	Mortality (48 hr, NOEC)	7.1-8.7	18-21	>9		135-217	<200	A,S,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i> straus	<24 hr old	Mortality (48 hr, LC50)	7.1-8.7	18-21	>9		135-217	226	A,S,M	P	Lewis and Valentine, 1981	Boric acid
Mosquito larvae	<i>Anopheles quadrimaculatus</i>		Mortality (48 hr, 92% mortality)						25	A	U	Fay, 1959	Boric acid
Mosquito larvae	3 species	through hatching	Mortality (LC97-LC99, through hatching)						43.7	A	?	U.S. EPA, 1975 In Eisler, 1990	Boric acid
Mosquito larvae	<i>Anopheles quadrimaculatus</i>		Mortality (25hr, 100% mortality)						125	A	U	Fay, 1959	Boric acid
Mosquito larvae	3 species	second instar	Mortality (48 hr, LC100)						524	A	?	U.S. EPA, 1975 In Eisler, 1990	Boric acid
Mosquito larvae	3 species	freshly hatched	Mortality (48 hr, LC100)						700	A	?	U.S. EPA, 1975 In Eisler, 1990	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
Mosquito larvae	3 species	third instar	Mortality (48 hr, LC100)						1,748	A	?	U.S. EPA, 1975 In Eisler, 1990	Boric acid
Mosquito larvae	3 species	pupae	Mortality (48 hr, LC100)						2,797	A	?	U.S. EPA, 1975 In Eisler, 1990	Boric acid
Midge	<i>Chironomus tentans</i>	third instar larval	Mortality (96 hr, LC50)		22-24			100	118	A	P*	MELP, 1996	Boric acid
Midge	<i>Chironomus tentans</i>	third instar larval	Mortality (96 hr, LC50)		22-24			250	137.7	A	P*	MELP, 1996	Boric acid
Midge	<i>Chironomus tentans</i>	third instar larval	Mortality (96 hr, LC50)		22-24			25	157.3	A	P*	MELP, 1996	Boric acid
Midge	<i>Chironomus tentans</i>	third instar larval	Mortality (48 hr, LC50)	7.5-7.9	22	5-8.4	85-90	120	242	A,S,M	P	MOE 2007 (unpublished data)	Boric acid
Midge	<i>Chironomus decorus</i>	larvae, fourth instar	Mortality (48 hr, LC50)	9.0-9.2	19.9-20.1	8.4-8.8		10.6-170	1,376	A,R,M	P	Maier and Knight, 1991	Sodium tetraborate
VERTEBRATES – Chronic													
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	(28 d, NOEC)	7.9	13.3	9.6	82	200	0.001	C,F,	S	Birge and Black, 1977	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	Mortality (28 d, LC1)	7.9	13.3	9.6	82	200	0.001	C,F,	S	Birge and Black, 1977	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	(28 d, LOEC)	7.9	13.3	9.6	82	200	0.01 ^b	C,F,	S	Birge and Black, 1977	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	freshly fertilized eggs	(32 d, NOEC)					200	0.01	C,F,	NA (Proprietary information)	Birge and Black, 1981 In Butterwick <i>et al.</i> , 1989	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>		Mortality (NOEC)						0.04	C	S	ANZECC, 2000	
Rainbow trout	<i>Oncorhynchus mykiss</i> ;	embryo larval	Mortality (28 d, LC1)	7.9	14	10.1		50	0.07	C,F,	S	Birge and Black, 1977	Borax

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
	<i>Salmo gairdneri</i>	stages											
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	Mortality (28 d, LC1)	7.8	13	10.3	82	200	0.07	C,F,	S	Birge and Black, 1977	Borax
Rainbow trout	<i>Oncorhynchus mykiss</i>		Mortality (32 d, LOEC)						~0.1	C	S	ANZECC, 2000	
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	freshly fertilized eggs	(32 d, LOEC)					200	0.1 ^b	C,F,	NA Proprietary information	Birge and Black, 1981 In Butterwick <i>et al.</i> , 1989	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	Mortality (28 d, LC1)	7.7	13.7	9.2		50	0.1	C,F,	S	Birge and Black, 1977	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Mortality at Hatching (32 d, LOEC)	7.4	13.2	9.8		197	0.1 ^b	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Teratogenesis at Hatching (32 d, LOEC)	7.4	13.2	9.8		197	0.1 ^b	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Mortality at 8-d Post-hatch (36 d, NOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	0.103	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Teratogenesis at 8-d Post-hatch (36 d, NOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	0.103	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	embryo-hatching stages	Mortality at Hatching (28 d, NOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	0.103	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo-larval stages	(28 d, NOEC)	7.7	13.7	9.2		50	0.11	C,F,	S	Birge and Black, 1977	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	early life stages	(36 d, NOEC)						0.75	C,F,	NA (Proprietary information)	Proctor and Gamble (unpublished) In Butterwick <i>et al.</i> , 1989	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo-larval stages	(28 d, NOEC)	7.9	14	10.1		50	0.96	C,F,	S	Birge and Black, 1977	Borax

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo-larval stages	(28 d, LOEC)	7.7	13.7	9.2		50	1 ^b	C,F,	S	Birge and Black, 1977	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	early life stages	(36 d, LOEC)						1	C,F,	NA (Proprietary information)	Proctor and Gamble (unpublished) In Butterwick <i>et al.</i> , 1989	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Mortality at 8-d Post-hatch (36 d, LOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	1.34	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Teratogenesis at 8-d Post-hatch (36 d, LOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	1.34	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-hatching stages	Mortality at Hatching (28 d, LOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	1.34	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-hatching stages	Teratogenesis at Hatching (28 d, NOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	1.34	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Mortality (larval survival) at 60-d Post-hatch (87 d, NOEC)	6.8-7.1	11-13		25-38	24-39	2.1	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Larval Growth at 60-d Post-hatch (87 d, NOEC)	6.8-7.1	11-13		25-38	24-39	2.1	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Embryo Survival at 60-d Post-hatch (87 d, NOEC)	6.8-7.1	11-13		25-38	24-39	2.1	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-larval stages	Embryo Viability at 60-d Post-hatch (87 d, NOEC)	6.8-7.1	11-13		25-38	24-39	2.1	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ;	embryo larval	(28 d, NOEC)	7.8	13	10.3	82	200	9.63	C,F,	S	Birge and Black, 1977	Borax

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
	<i>Salmo gairdneri</i>	stages											
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	(28 d, LOEC)	7.9	14	10.1		50	9.7	C,F,	S	Birge and Black, 1977	Borax
Rainbow Trout	<i>Oncorhynchus mykiss</i>	embryo-hatching stages	Teratogenesis at Hatching (28 d, LOEC)	7.5-7.9	13.4-14	9.7-10.3	47-61	178-198	11.46	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	early life stages	(60 d, LOEC)	6.5-7.5				27	>17	C,F,	NA (Proprietary information)	Proctor and Gamble (unpublished) In Butterwick <i>et al.</i> , 1989	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	20 d embryo-larval stages	Larval Growth at 60-d Post-hatch (67 d, NOEC)	6.8-7.1	11-13		25-38	24-39	18	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	20 d embryo-larval stages	Embryo Survival at 60-d Post-hatch (67 d, NOEC)	6.8-7.1	11-13		25-38	24-39	18	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	20 d embryo-larval stages	Larval Survival at 60-d Post-hatch (67 d, NOEC)	6.8-7.1	11-13		25-38	24-39	18	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	Mortality (28 d, LC50)	7.9	14	10.1		50	27	C,F,	S	Birge and Black, 1977	Borax
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	(28 d, LOEC)	7.8	13	10.3	82	200	49.7	C,F,	S	Birge and Black, 1977	Borax
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	Mortality (28 d, LC50)	7.8	13	10.3	82	200	54	C,F,	S	Birge and Black, 1977	Borax
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	Mortality (28 d, LC50)	7.9	13.3	9.6	82	200	79	C,F,	S	Birge and Black, 1977	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>	embryo larval stages	Mortality (28 d, LC50)	7.7	13.7	9.2		50	100	C,F,	S	Birge and Black, 1977	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
Rainbow trout	<i>Oncorhynchus mykiss</i>	embryo (fertilization to 2 weeks post-hatch)	Embryonic Mortality (6 wk, LOEC, 85 to 95% mortality)		12.5				108 (10 mmol B/L) ^a	C,M	S	Rowe <i>et al.</i> , 1998	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>	embryo larval stages	Mortality (32 d, LC50)	7.4	13.2	9.8		197	138	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Rainbow trout	<i>Oncorhynchus mykiss</i>		Mortality (32 d, LC50)						~138	C	S	ANZECC, 2000	
Rainbow trout	<i>Oncorhynchus mykiss</i> ; <i>Salmo gairdneri</i>		Darkening of skin, immobilization and loss of equilibrium						874	C	? (Original publication in French only)	Wurtz, 1945 In Butterwick <i>et al.</i> , 1989	Boric acid
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC1)	7.6	24.8	7.5	82	200	0.2	C,F,	S	Birge and Black, 1977	Boric acid
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC1)	7.9	24.8	7.4		50	0.6	C,F,	S	Birge and Black, 1977	Boric acid
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC1)	8.1	27	7.5	82	200	0.9	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC1)	8.3	27	7.5		50	1.4	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC50)	8.3	27	7.5		50	6.5	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius auratus</i>	embryo larval stages	(7 d, NOEC)	7.6	24.8	7.5	82	200	6.8	C,F,	S	Birge and Black, 1977	Boric acid
Goldfish	<i>Carassius auratus</i>	embryo larval stages	(7 d, LOEC)	7.6	24.8	7.5	82	200	8.33	C,F,	S	Birge and Black, 1977	Boric acid
Goldfish	<i>Carassius auratus</i>	embryo larval stages	(7 d, NOEC)	8.1	27	7.5	82	200	8.53	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius</i>	embryo	(7 d, NOEC)	7.9	24.8	7.4		50	9.2	C,F,	S	Birge and	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
	<i>auratus</i>	larval stages										Black, 1977	
Goldfish	<i>Carassius auratus</i>	embryo larval stages	(7 d, LOEC)	7.9	24.8	7.4		50	22.5	C,F,	S	Birge and Black, 1977	Boric acid
Goldfish	<i>Carassius auratus</i>	embryo larval stages	(7 d, NOEC)	8.3	27	7.5		50	26.5	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius auratus</i>	embryo larval stages	(7 d, LOEC)	8.1	27	7.5	82	200	27.33	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC50)	7.9	24.8	7.4		50	46	C,F,	S	Birge and Black, 1977	Boric acid
Goldfish	<i>Carassius auratus</i>	embryo larval stages	(7 d, LOEC)	8.3	27	7.5		50	48.75	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC50)	8.1	27	7.5	82	200	59	C,F,	S	Birge and Black, 1977	Borax
Goldfish	<i>Carassius auratus</i>	embryo larval stages	Mortality (7 d, LC50)	7.6	24.8	7.5	82	200	75	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC1)	7.6	24.7	7.6	82	200	0.2	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	(9 d, NOEC)	8.2	29.4	6.5	82	200	0.49	C,F,	S	Birge and Black, 1977	Borax
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC1)	7.5	25	7.3		50	0.5	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	(9 d, NOEC)	7.6	24.7	7.6	82	200	0.75	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	(9 d, LOEC)	7.6	24.7	7.6	82	200	1	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval	(9 d, NOEC)	7.5	25	7.3		50	1.01	C,F,	S	Birge and Black, 1977	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
		stages											
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	(9 d, LOEC)	8.2	29.4	6.5	82	200	1.04	C,F,	S	Birge and Black, 1977	Borax
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC1)	8.2	29.4	6.5	82	200	1.7	C,F,	S	Birge and Black, 1977	Borax
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	(9 d, LOEC)	7.5	25	7.3		50	5.42	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC1)	8.5	29.4	6.4		50	5.5	C,F,	S	Birge and Black, 1977	Borax
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	(9 d, NOEC)	8.5	29.4	6.4		50	9	C,F,	S	Birge and Black, 1977	Borax
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC50)	7.6	24.7	7.6	82	200	22	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	(9 d, LOEC)	8.5	29.4	6.4		50	25.9	C,F,	S	Birge and Black, 1977	Borax
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC50)	8.2	29.4	6.5	82	200	71	C,F,	S	Birge and Black, 1977	Borax
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC50)	7.5	25	7.3		50	155	C,F,	S	Birge and Black, 1977	Boric acid
Channel catfish	<i>Ictalurus punctatus</i>	embryo larval stages	Mortality (9 d, LC50)	8.5	29.4	6.4		50	155	C,F,	S	Birge and Black, 1977	Borax
Largemouth bass	<i>Micropterus salmoides</i>	embryo-larval stages	Mortality at 8-d Post-hatch (11 d, NOEC)	7.5	20	8.4		204	1.39	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Largemouth bass	<i>Micropterus salmoides</i>	freshly fertilized eggs	(11 d, NOEC)					200	1.39	C,F,	NA (Proprietary information)	Birge and Black, 1981 In Butterwick <i>et al.</i> , 1989	Boric acid
Largemouth bass	<i>Micropterus salmoides</i>	embryo-larval	Mortality at 8-d Post-hatch (11	7.5	20	8.4		204	12.17	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
		stages	d, LOEC)										
Largemouth bass	<i>Micropterus salmoides</i>	freshly fertilized eggs	(11 d, LOEC)					200	12.17	C,F,	NA (Proprietary information)	Birge and Black, 1981 In Butterwick <i>et al.</i> , 1989	Boric acid
Largemouth bass	<i>Micropterus salmoides</i>	embryo-larval stages	Mortality (11 d, LC50)	7.5	20	8.4		204	92	C,F,M	P	Black <i>et al.</i> , 1993	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	Mortality (7.5 d, LC1)	8.4	25.3	7.8	82	200	3	C,F,	S	Birge and Black, 1977	Borax
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	Mortality (7.5 d, LC1)	8.3	25.3	7.7		50	5	C,F,	S	Birge and Black, 1977	Borax
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d NOEC)	8.4	25.3	7.8	82	200	7.04	C,F,	S	Birge and Black, 1977	Borax
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d NOEC)	8.3	25.3	7.7		50	7.04	C,F,	S	Birge and Black, 1977	Borax
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d, LOEC)	8.3	25.3	7.7		50	9.6	C,F,	S	Birge and Black, 1977	Borax
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d, LOEC)	8.4	25.3	7.8	82	200	10.5	C,F,	S	Birge and Black, 1977	Borax
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	Mortality (7.5 d, LC1)	7.7	25	7.7		50	13	C,F,	S	Birge and Black, 1977	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	Mortality (7.5 d, LC1)	7.7	25	7.8	82	200	22	C,F,	S	Birge and Black, 1977	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d NOEC)	7.7	25	7.7		50	32.5	C,F,	S	Birge and Black, 1977	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d NOEC)	7.7	25	7.8	82	200	45.7	C,F,	S	Birge and Black, 1977	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval	Mortality (7.5 d, LC50)	8.3	25.3	7.7		50	47	C,F,	S	Birge and Black, 1977	Borax

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
		stages											
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d, LOEC)	7.7	25	7.7		50	47.5	C,F,	S	Birge and Black, 1977	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	Mortality (7.5 d, LC50)	8.4	25.3	7.8	82	200	54	C,F,	S	Birge and Black, 1977	Borax
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	(7 d, LOEC)	7.7	25	7.8	82	200	86	C,F,	S	Birge and Black, 1977	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	Mortality (7.5 d, LC50)	7.7	25	7.7		50	130	C,F,	S	Birge and Black, 1977	Boric acid
Leopard Frog	<i>Rana pipiens</i>	embryo-larval stages	Mortality (7.5 d, LC50)	7.7	25	7.8	82	200	135	C,F,	S	Birge and Black, 1977	Boric acid
Zebrafish	<i>Brachydanio rerio</i>		mortality, growth, condition (34-d MATC)						10.04		?	Hooftman et al, 2000 cited in U.S. Borax Inc MSDS	
Fathead minnow	<i>Pimeohales promelas</i>	eggs and fry	Growth Reduction (30 d, NOEC)	7.1-7.9	25		33-38	38-46	14	C,F,	S	Proctor and Gamble (unpublished) In Moss and Nagpal, 2003	Boric acid
Fathead minnow	<i>Pimephales promelas</i>	Larval	Growth Inhibition (7d IC25)	7.5-7.9	18-22	5-8.4	85-90	120	20.6 (8.5-26.5)	C,R,M	P	MOE 2007 (unpublished data)	
Fathead minnow	<i>Pimephales promelas</i>	eggs and fry	Growth Reduction (30 d, LOEC)	7.1-7.9	25		33-38	38-46	24	C,F,	S	Proctor and Gamble (unpublished) In Moss and Nagpal, 2003	Boric acid
Fathead minnow	<i>Pimephales promelas</i>	eggs and fry	Reduced Fry Survival (60 d, NOEC)	7.1-7.9	25		33-38	38-46	24	C,F,	S	Proctor and Gamble (unpublished) In Moss and Nagpal, 2003	Boric acid
Fathead minnow	<i>Pimeohales promelas</i>	eggs and fry	Reduced Fry Survival	7.1-7.9	25		33-38	38-46	88	C,F,	S)	Proctor and Gamble	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
			(60 d, LOEC)								(unpublished) In Moss and Nagpal, 2003		
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	Mortality (7 d, LC1)	7.6	23.7	6.8	85	200	5	C,F,	S	Birge and Black, 1977	Boric acid
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	(7 d, NOEC)	7.6	23.7	6.8	82	200	22.3	C,F,	S	Birge and Black, 1977	Boric acid
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	Mortality (7.5 d, LC1)	7.6	23.7	6.8		50	25	C,F,	S	Birge and Black, 1977	Boric acid
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	(7 d, NOEC)	7.6	23.7	6.8		50	48.7	C,F,	S	Birge and Black, 1977	Boric acid
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	(7 d, LOEC)	7.6	23.7	6.8	82	200	53.5	C,F,	S	Birge and Black, 1977	Boric acid
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	(7 d, LOEC)	7.6	23.7	6.8		50	96	C,F,	S	Birge and Black, 1977	Boric acid
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	Mortality (7d, LC50)	7.6	23.7	6.8	82	200	123	C,F,	S	Birge and Black, 1977	Boric acid
Fowler's toad	<i>Bufo fowleri</i>	embryo-larval stages	Mortality (7.5 d, LC50)	7.6	23.7	6.8		50	145	C,F,	S	Birge and Black, 1977	Boric acid
Wood frog	<i>Rana sylvatica</i>	blastula lifestage eggs	Proportion of Deformed Hatchlings (13-23 d, LOEC, 51% increase in deformities compared to controls)	6.5	10				50	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
Wood frog	<i>Rana sylvatica</i>	blastula lifestage eggs	Proportion of Eggs Hatched (13-23 d, NOEC)	6.5	10				100	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
Wood frog	<i>Rana sylvatica</i>	blastula lifestage eggs	Proportion of Deformed Hatchlings (13-23 d, 77% increase in deformities compared to controls)	6.5	10				100	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
Jefferson salamander	<i>Ambystoma jeffersonianum</i>	blastula lifestage eggs	Proportion of Deformed Hatchlings (17-25 d, LOEC, 7% increase in deformities compared to controls)	6.5	10				50	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
Jefferson salamander	<i>Ambystoma jeffersonianum</i>	blastula lifestage eggs	Proportion of Eggs Hatched (17-25 d, NOEC)	6.5	10				100	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
Jefferson salamander	<i>Ambystoma jeffersonianum</i>	blastula lifestage eggs	Proportion of Deformed Hatchlings (17-25 d, 13% increase in deformities compared to controls)	6.5	10				100	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
Spotted salamander	<i>Ambystoma maculatum</i>	blastula lifestage eggs	Proportion of Deformed Hatchlings (38-44 d, LOEC, 18% increase in deformities compared to controls)	6.5	10				50	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
Spotted salamander	<i>Ambystoma maculatum</i>	(blastula lifestage)	Proportion of Eggs Hatched	6.5	10				100	C,S,M	S	Laposata and Dunson,	Sodium tetraborate

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
		eggs	(38-44 d, NOEC)								1998		
Spotted salamander	<i>Ambystoma maculatum</i>	blastula lifestage eggs	Proportion of Deformed Hatchlings (38-44 d, 80% increase in deformities compared to controls)	6.5	10				100	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
American toad	<i>Bufo americanus</i>	blastula lifestage eggs	Proportion of Eggs Hatching (15-23 d, LOEC, 43% decrease compared to controls)	6.5	10				50	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
American toad	<i>Bufo americanus</i>	blastula lifestage eggs	Proportion of Eggs Hatching (15-23 d, 71% decrease compared to controls)	6.5	10				100	C,S,M	S	Laposata and Dunson, 1998	Sodium tetraborate
Coho salmon	<i>Onchorhynchus kisutch</i>	alevins and fry	Mortality (23 d, LC50)	7	11				93	C,R	S	Davis and Mason, 1973	Sodium metaborate
Coho salmon	<i>Onchorhynchus kisutch</i>	alevins 0.19-0.7 g	Mortality (283 hr, LC50)		11			47	113 (104-123 95% CLs)	C,R	S	Thompson <i>et al.</i> , 1976	Sodium metaborate
INVERTEBRATES – Chronic													
Water Flea	<i>Daphnia magna</i>		Immobilization (threshold concentration)		25				<0.38	C	?	McKee and Wolf, 1963 In Butterwick <i>et al.</i> , 1989	Sodium perborate
Water Flea	<i>Daphnia magna</i>		Growth (21d MATC)						4.665	C	S	ANZECC, 2000	
Water Flea	<i>Daphnia magna</i>		Reproduction						6	C	S	ANZECC,	

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
			(NOEC)								2000		
Water Flea	<i>Daphnia magna</i> straus		Mean Brood Sizes (21 d, NOEC)	7.1-8.7	18-21	>9		135-217	6	C,R,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i>		(21 d, NOEC)	8.0-8.2	19.5-20.5	7.3-8.0	53-63	141-155	6.4	C,S,R,M	S	Gersich, 1984	Boric acid
Water Flea	<i>Daphnia magna</i>		Growth (21 d, MATC)	8.0-8.2	19.5-20.5	7.3-8.0	53-63	141-155	~9.3	C,S,R,M	S	Gersich, 1984	Boric acid
Water Flea	<i>Daphnia magna</i>	(<24 hr old)	(NOEC)					250	12.4	C,S	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i> straus		Mean Brood Sizes (21 d, LOEC)	7.1-8.7	18-21	>9		135-217	13	C,R,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i>	(<24 hr old)	(NOEC)					100	13.1	C,S	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i> straus		(21 d, LOEC)	8.0-8.2	19.5-20.5	7.3-8.0	53-63	141-155	13.6	C,S,R,M	S	Gersich, 1984	Boric acid
Water Flea	<i>Daphnia magna</i> straus		Reproduction (LOEC, mean number of broods per daphnid, mean total number of young per daphnid, mean brood size per daphnid, and mean size)	8.0-8.2	19.5-20.5	7.3-8.0	53-63	141-155	14	C,S,R,M	S	Gersich, 1984	Boric acid
Water Flea	<i>Daphnia magna</i>	(<24 hr old)	(LOEC)					100	25.4	C,S	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i>	(<24 hr old)	(LOEC)					250	26.4	C,S	P*	MELP, 1996	Boric acid
Water Flea	<i>Daphnia magna</i> straus		Mean Length (21 d, NOEC)	7.1-8.7	18-21	>9		135-217	27	C,R,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i> straus		Mean Brood Sizes (21 d, 31% decrease compared to	7.1-8.7	18-21	>9		135-217	27	C,R,M	P	Lewis and Valentine, 1981	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
			controls)										
Water Flea	<i>Daphnia magna</i>		Immobilization (threshold concentration)						<27.2	C	?	McKee and Wolf, 1963 In Butterwick <i>et al.</i> , 1989	Sodium tetraborate
Water Flea	<i>Daphnia magna</i> straus		Mortality (21 d, LC50)	7.3-8.0	20			150	52.2	C,S,R,M	P	Gersich, 1984	Boric acid
Water Flea	<i>Daphnia magna</i> straus		Mean Length (21 d, LOEC)	7.1-8.7	18-21	>9		135-217	53	C,R,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i> straus		Number of Offspring Produced (21 d, 70% decrease compared to controls)	7.1-8.7	18-21	>9		135-217	53	C,R,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i> straus		Mean Brood Sizes (21 d, 50% decrease compared to controls)	7.1-8.7	18-21	>9		135-217	53	C,R,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i> straus	adult	Mortality (21 d, LC50)	7.1-8.7	18-21	>9		135-217	53.2 (44.1-64.5)	C,R,M	P	Lewis and Valentine, 1981	Boric acid
Water Flea	<i>Daphnia magna</i>		Mortality (21 d, LC50)						~266	C	S	ANZECC, 2000	
Water Flea	<i>Ceriodaphnia dubia</i>		Growth, reproduction (14-d MATC)						13.4		S	Hickey, 1989	
Amphipod	<i>Hyalella azteca</i>	1-11 d	Mortality (7 d, LC50)	7.23-8.83	24-25		14	18	2.935	A,S,M	S	Borgmann <i>et al.</i> , 2005	Boron
Amphipod	<i>Hyalella azteca</i>	1-11 d	Mortality (7 d, LC50)	6.44-8.52	24-25		84	124	>3.15	A,S,U	S	Borgmann <i>et al.</i> , 2005	Boron
Unknown - Protozoa???	<i>Opercularia bimarginata</i>		Growth, reproduction (72-h NOEC)						10		S	Guhl, 1992b, In Dyer 2001	
Protozoa	<i>Entosiphon sulcatum</i>		Growth (72-h NOEC)						15		S	Guhl 1992b In Dyer	

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
											2001		
Protozoa	<i>Paramecium caudatum</i>		Growth, reproduction (72-h NOEC)						20		S	Guhl 1992b, In Dyer 2001	
Midge	<i>Chironomus decorus</i>	larvae, fourth instar	Growth Rate (96 hr, NOEC, inhibited growth rate)	9.0-9.2	19.9-20.1	8.4-8.8		10.6-170	<20	C,R,M	P	Maier and Knight, 1991	Sodium tetraborate
Midge	<i>Chironomus decorus</i>	larvae, fourth instar	Growth Rate (96 hr, LOEC, significantly inhibited growth rate)	9.0-9.2	19.9-20.1	8.4-8.8		10.6-170	20	C,R,M	P	Maier and Knight, 1991	Sodium tetraborate
<u>PLANTS – Chronic</u>													
Blue Green Algae	<i>Anacystis nidulans</i>		Growth (NOEC)						0.01-4.0	C	S	Martinez <i>et al.</i> , 1986	Boric acid
Blue Green Algae	<i>Anacystis nidulans</i>		Growth or Organic Constituents (NOEC)						50	C	S	Martinez <i>et al.</i> , 1986	Boric acid
Blue Green Algae	<i>Anacystis nidulans</i>		Significantly decreased growth and chlorophyll content						75	C	S	Martinez <i>et al.</i> , 1986	Boric acid
Blue Green Algae	<i>Anacystis nidulans</i>		Decreased chlorophyll content and photosynthesis inhibition within 72 hrs						100	C	S	Martinez <i>et al.</i> , 1986	Boric acid
Blue Green Algae	<i>Anacystis nidulans</i>		Decreased protein content causing inhibition in nitrate uptake and nitrate reductase						100	C	S	Martinez <i>et al.</i> , 1986	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
			activity										
Green Alga	<i>Chlorella pyrenoidosa</i>		Population Growth (14 d, NOEC)						0.4	C	S	ANZECC, 2000	
Green Alga	<i>Chlorella pyrenoidosa</i>		Growth or Cell Composition (NOEC)						10	C	U	Fernandez <i>et al.</i> , 1984	
Green Alga	<i>Chlorella pyrenoidosa</i>		Decreased algal growth; increase in protein and nucleic acid synthesis						≥25	C	? Publication in Spanish	Sanchez <i>et al.</i> , 1982 In Maier and Knight, 1991	Boron
Green Alga	<i>Chlorella pyrenoidosa</i>		Altered cell division and amino acid activity after 72 hr; reversible photosynthesis inhibition; giant cells formed with increased nitrate and protein						50-100	C	U	Maeso <i>et al.</i> , 1985	
Green Alga	<i>Chlorella pyrenoidosa</i>		Cell Division and Biomass Synthesis (72 hr, totally inhibitory)						>100	C	U	Maeso <i>et al.</i> , 1985	
Duckweed	<i>Spirodella polyrrhiza</i>		FronD Production (10 d, NOEC)	5.2-5.8	24-26				0.9	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
Duckweed	<i>Spirodella polyrrhiza</i>		FronD Production (10 d, LOEC)	5.2-5.8	24-26				3.5	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
Duckweed	<i>Spirodella polyrrhiza</i>		Growth Rate (10d, NOEC)	5.2-5.8	24-26				6.1	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
Duckweed	<i>Spirodella polyrrhiza</i>		Reduced Growth Rate	5.2-5.8	24-26				11.7	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
			(10d, EC50)										
Duckweed	<i>Spirodella polyrrhiza</i>		Reduced Frond Production (10 d, EC50)	5.2-5.8	24-26				14.3	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
Duckweed	<i>Spirodella polyrrhiza</i>		Abnormal Frond Occurrence (chlorotic, necrotic, dead) (10 d, EC50)	5.2-5.8	24-26				17.7	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
Duckweed	<i>Spirodella polyrrhiza</i>		Growth Rate (10d, LOEC)	5.2-5.8	24-26				18.9	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
Duckweed	<i>Spirodella polyrrhiza</i>		Abnormal Fronds (chlorotic, necrotic, dead) (10 d, NOEC)	5.2-5.8	24-26				18.9	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
Duckweed	<i>Spirodella polyrrhiza</i>		Abnormal Fronds (chlorotic, necrotic, dead) (10 d, LOEC)	5.2-5.8	24-26				22.4	C,R,M	S	Davis <i>et al.</i> , 2002	Boric acid
American waterweed	<i>Elodea canadensis</i>		Growth inhibition						1	C	S*	Nobel <i>et al.</i> , 1983 In Maier and Knight, 1991	Boric acid
American waterweed	<i>Elodea canadensis</i>		(NOEC)						1	C	S	ANZECC, 2000	
American waterweed	<i>Elodea canadensis</i>		Mortality (21 d, LC50)						~5.0	C	S	ANZECC, 2000	
Stream water-crowfoot	<i>Ranunculus penicillatus</i>		Growth inhibition						1	C	S*	Nobel <i>et al.</i> , 1983 In Maier and Knight, 1991	
Water milfoil	<i>Myriophyllum alterniflorum</i>		Growth inhibition						2	C	S*	Nobel <i>et al.</i> , 1983 In Maier and	

Aquatic Toxicity Table for Boron

COMMON NAME	SPECIES NAME	LIFE STAGE	RESPONSE	TEST CONDITIONS					EFFECT CONC (mg/L)	DATA CODES KEY(1)	DATA QUALITY KEY(2)	DATA REF	Salt Used
				pH	TEMP (°C)	DO (mg/L)	ALK. (mg/L)	HARD (mg/L)					
											Knight, 1991		
Common reed	<i>Phragmites australis</i>		growth (4-month, NOEC)						< 4		S	Bergmann 1995	
Duckweed	<i>Lemna minor</i>		Decreased fresh weight per plant						5	C	S	Frick, 1985 In Maier and Knight, 1991	Boric acid
Duckweed	<i>Lemna minor</i>		Growth (7-d NOEC)						60		S	Wang, 1986	
Duckweed	<i>Lemna minor</i>		Growth inhibition	5					100	C	S	Frick, 1985 In Maier and Knight, 1991	Boric acid
Chlorella algae	<i>Chlorella vulgaris</i>		Population Growth (NOEC)						5.2	C	S	ANZECC, 2000	
Green Alga	<i>Selenastrum capricornutum</i>	4-7 d old	NOEC (72 hr)		22-26				<12.3	C,S	P*	MELP, 1997	
Green Alga	<i>Selenastrum capricornutum</i>	4-7 d old	LOEC (72 hr)		22-26				12.3	C,S	P*	MELP, 1997	
Green Alga	<i>Scenedesmus subpicatus</i>		Growth (96-h EC 10)						30		S	Guhl 1992, Dyer 2001	
Spiked or Eurasian watermillfoil	<i>Myriophyllum spicatum</i>		(32 d, EC50)						~171	C	S	ANZECC, 2000	
Spiked or Eurasian watermillfoil	<i>Myriophyllum spicatum</i>		(NOEC)						34.2	C	S	ANZECC, 2000	
Spiked or Eurasian watermillfoil	<i>Myriophyllum spicatum</i>		Root Growth (32 d, 50% inhibition of roots weight)						40.3	C	U	Stanley, 1974	

* Indicates values entered into Table

Assign 3 data codes, one from each of the following rows:		
A-acute	C-chronic	
S-static	R-static/renewal	F-flowthrough
U-unmeasured nominal concentration	M-measured concentration	

P Primary study	S Secondary Study	U Unacceptable Study
-----------------	-------------------	----------------------

TLm- (median tolerance limit) This is the concentration at which acute toxicity ceases, usually taken as the concentration at which 50% of the population of test organisms can live for an indefinite time. This endpoint is now commonly known as an LC50.

^a Converted using a molecular weight of 10.812 g/mol

^b Chronic toxicity data points were not used for iPWQG development as these data points were considered outliers as other scientists and studies have not been able to reproduce these low values using similar conditions and species (Moss and Nagpal, 2003)

P* Classified as primary by the BC MOE

S* Classified as secondary by the BC MOE as test conditions were not adequately reported

? Not classifiable

NA Report was not available at all, or not available in English